


OBSERVATION OF SINGLE TOP AT CDF

EPS-HEP2009 • Kraków, Poland • July 16-22, 2009

Bruno Casal Laraña



Instituto de Física de Cantabria
(CSIC-Univ. de Cantabria)

On behalf of the CDF Collaboration





THE TEVATRON AT FERMILAB

- $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV
- Peak luminosity $\sim 360 \times 10^{30} \text{cm}^{-2}\text{s}^{-1}$
- Tevatron delivered $\sim 6.9 \text{fb}^{-1}$
- CDF collected $\sim 5.7 \text{fb}^{-1}$
 - Analysis shown here use 3.2fb^{-1}
- Tevatron is performing extremely well
 - Expected $>10 \text{fb}^{-1}$ by end of 2010
 - Possibly run in 2011 and beyond



THE CDF II DETECTOR

- Multipurpose detector at Tevatron collecting data efficiently
 - Large acceptance and good ID for leptons
 - Tracking and EM calorimeter
 - Muon systems
 - Good calorimetry for jet energy resolution
 - Silicon detectors for b-jet tagging



MOTIVATION

- Single top quark produced via electroweak interaction



- $\sigma_{SM}(t\text{-channel}) = 1.98 \pm 0.21 \text{ pb}$
- $\sigma_{SM}(s\text{-channel}) = 0.88 \pm 0.07 \text{ pb}$
- $\sigma_{SM}(t\bar{t}) = 6.7 \pm 0.8 \text{ pb}$ (via strong interaction)
- [Z. Sullivan, Phys. Rev. D70, 114012 (2004)] ($m_t = 175 \text{ GeV}$)

- Test of the Standard Model

- Direct measurement of $|V_{tb}|$
- Top quark properties: polarization, spin, W helicity,...
- Same final state as WH


- Sensitive to new physics

- Search for W' , H^+ (s-channel signature)
- Search for FCNC,...




THE CHALLENGE

- Rare process at the Tevatron
 - $S : B \sim 1 : 10^9$ before trigger
- Not very distinct signature with many sources of background
 - $S : B \sim 1 : 20$ after final selection
 - No “golden” variable
- Large systematics uncertainties
- Sophisticated analysis methods needed
- Current approaches at CDF: Likelihood Function, Matrix Element, Neural Network, Boosted Decision Trees
 - $S : B > 1 : 1$ in most significant bins




SINGLE TOP SELECTION AT CDF

- Top decays most of the times to Wb
- $W + 2$ or 3 jets with $E_T > 20$ GeV
- One lepton (electron or muon) with $p_T > 20$ GeV
 - Mainly from standard high p_T lepton triggers
 - Extended muon coverage from missing E_T dedicated triggers (30% gain in signal acceptance)
- Missing transverse energy from neutrino, $\cancel{E}_T > 25$ GeV
- Veto “non- W ”, Z , dilepton, conversion, cosmics
- At least one b -tagged jet (displaced secondary vertex algorithm)
- Main backgrounds: W +Heavy Flavor, W +Mistags, $t\bar{t}$,...







Process	Number of Events in 3.2 fb ⁻¹	
	W + 2 jets	W + 3 jets
s-channel	58.1 ± 8.4	19.2 ± 2.8
t-channel	87.6 ± 13.0	26.2 ± 3.9
$Wb\bar{b}$	656.9 ± 198.0	201.3 ± 60.8
$Wc\bar{c}$	292.2 ± 90.1	98.1 ± 30.2
Wcj	250.4 ± 77.2	52.1 ± 16.0
Mistags	501.3 ± 69.6	151.9 ± 21.4
non- W	89.6 ± 35.8	35.1 ± 14.0
WW	58.5 ± 6.6	21.2 ± 2.4
WZ	28.9 ± 2.4	8.5 ± 0.7
ZZ	0.9 ± 0.1	0.4 ± 0.0
$Z + jets$	36.5 ± 5.6	15.6 ± 2.4
$t\bar{t}$ dilepton	69.2 ± 10.0	60.2 ± 8.7
$t\bar{t}$ non-dilepton	134.9 ± 19.6	421.8 ± 61.1
Total signal	145.7 ± 21.4	45.4 ± 6.7
Total prediction	2265.0 ± 375.4	1111.5 ± 129.5
Observed in data	2229	1086

MONTE CARLO MODELING







MONTE CARLO MODELING




CHECKING HUNDRED OF PLOTS!!



- Data versus MC agreement
- Input Variables and more
- Control and signal regions




MULTIVARIATE ANALYSES





- Once the background estimate is well understood we are ready to perform a multivariate analysis
 - Neural Network (NN)**
 - Boosted Decision Tree (BDT)
 - Matrix Element Method (ME)
 - Likelihood Function (LF)
 - Dedicated t- and s-channel searches
- Discriminant outputs are thoroughly cross-checked in sideband regions before unblinding the signal region




MULTIVARIATE ANALYSES



- Once the background estimate is well understood we are ready to perform a multivariate analysis
 - Neural Network (NN)
 - Boosted Decision Tree (BDT)**
 - Matrix Element Method (ME)
 - Likelihood Function (LF)
 - Dedicated t- and s-channel searches
- Discriminant outputs are thoroughly cross-checked in sideband regions before unblinding the signal region




MULTIVARIATE ANALYSES




- Once the background estimate is well understood we are ready to perform a multivariate analysis
 - Neural Network (NN)
 - Boosted Decision Tree (BDT)
 - Matrix Element Method (ME)**
 - Likelihood Function (LF)
 - Dedicated t- and s-channel searches
- Discriminant outputs are thoroughly cross-checked in sideband regions before unblinding the signal region

$$EPD = \frac{b \cdot P_{sig}}{b \cdot P_{sig} + b \cdot P_{b-bkg} + (1-b) \cdot P_{nonb-bkg}}$$




MULTIVARIATE ANALYSES




- Once the background estimate is well understood we are ready to perform a multivariate analysis
 - Neural Network (NN)
 - Boosted Decision Tree (BDT)
 - Matrix Element Method (ME)
 - Likelihood Function (LF)**
 - Dedicated **t**- and **s**-channel searches

$$L = \frac{\prod_{i=1}^{n_{\text{var}}} p_i^{\text{sig}}(x_i)}{\prod_{i=1}^{n_{\text{var}}} p_i^{\text{sig}}(x_i) + \prod_{i=1}^{n_{\text{var}}} p_i^{\text{bkg}}(x_i)}$$

- Discriminant outputs are thoroughly cross-checked in sideband regions before unblinding the signal region




MULTIVARIATE ANALYSES




- Once the background estimate is well understood we are ready to perform a multivariate analysis
 - Neural Network (NN)
 - Boosted Decision Tree (BDT)
 - Matrix Element Method (ME)
 - Likelihood Function (LF)**
 - Dedicated t- and **s-channel** searches

$$L = \frac{\prod_{i=1}^{n_{\text{var}}} p_i^{\text{sig}}(x_i)}{\prod_{i=1}^{n_{\text{var}}} p_i^{\text{sig}}(x_i) + \prod_{i=1}^{n_{\text{var}}} p_i^{\text{bkg}}(x_i)}$$

- Discriminant outputs are thoroughly cross-checked in sideband regions before unblinding the signal region




MULTIVARIATE ANALYSES



- Once the background estimate is well understood we are ready to perform a multivariate analysis
 - Neural Network (NN)
 - Boosted Decision Tree (BDT)
 - Matrix Element Method (ME)
 - Likelihood Function (LF)
 - Dedicated t- and s-channel searches
- Discriminant outputs are thoroughly cross-checked in sideband regions before unblinding the signal region

BINNED LIKELIHOOD FIT AND SYSTEMATICS


- A binned maximum likelihood fit of the templates to the data is performed to extract the cross-section
- Systematic uncertainties are taken into account as nuisance parameters
 - Rate syst. give freedom to the fit templates to move vertically only
 - Shape syst. allow templates to slide horizontally (bin by bin)
 - MC statistics in each bin treated as a source of systematic uncertainty



Systematic Uncertainty	Rate	Shape
Jet Energy Scale	0...10%	✓
Initial + Final State Radiation	0...15%	✓
Parton Distribution Functions	2...3%	✓
Monte Carlo Generator	1...5%	
Event Detection Efficiency	0...9%	
Luminosity	6%	
Neural Net B-tagger		✓
Mistag Model		✓
Q ² scale in ALPGEN MC		✓
Input variable mismodeling		✓
Wbb+Wcc normalization	30%	
Wc normalization	30%	
Mistag normalization	17...29%	
ttbar normalization & m _{top}	23%	✓

SIGNIFICANCE AND RESULTS

- P-value is determined throwing pseudo experiments (PE) for the test hypothesis (S+B) and for the background-only hypothesis (B)
 - Fluctuate all rate and shape systematic uncertainties in PE
 - Test Statistic: $-2 \ln(Q)$, $Q = \mathcal{L}(S+B)/\mathcal{L}(B)$
 - Exp. p-value: fraction of B PE with a value smaller than the median of S+B PE
 - Obs. p-value: fraction of B PE with a value smaller than the measured one





CDF Preliminary, $L = 3.2 \text{ fb}^{-1}$			
Analysis	Exp. sig.	Obs. sig.	Cross sect. (pb)
NN	5.2σ	3.5σ	$1.8^{+0.6}_{-0.6}$
BDT	5.2σ	3.5σ	$2.1^{+0.7}_{-0.6}$
ME	4.9σ	4.3σ	$2.5^{+0.7}_{-0.6}$
LF	4.0σ	2.4σ	$1.6^{+0.8}_{-0.7}$
LFS (s-chan)	1.1σ	2.0σ	$1.5^{+0.9}_{-0.8}$


MET + JETS ANALYSIS (MJ)

- Another single top search in the orthogonal E_T +jets sample
 - Increase acceptance by $\sim 30\%$: events where lepton was not reconstructed and hadronic taus
 - Neural Network to suppress huge instrumental background
 - Final Neural Network to discriminate signal from background
 - Same method to extract signal cross-section and significance


CDF Preliminary, Luminosity = 2.1 fb^{-1}			
Analysis	Exp. sig.	Obs. sig.	Cross section (pb)
E_T +jets	1.4	2.1	$4.9^{+2.5}_{-2.2}$



COMBINATION TECHNIQUE





- Combine individual analysis into one “Super Discriminant”
 - Use discriminant outputs (NN, BDT, ME, LF, LFS) as input to NN
 - Candidate networks compete against each other
 - Optimization of network topology, weights, output histogram binning
 - Final network chosen based on best expected p-value
 - Gained 13% over best individual analysis




COMBINATION RESULTS

- Perform a simultaneous cross section fit over the two orthogonal analysis (SD + MJ)
- Expected p-value: $> 5.9\sigma$
- Observed p-value: **5.0σ**
- Measured cross-section: $\sigma = 2.3^{+0.6}_{-0.5} \text{ pb}$
- Extract $|V_{tb}| = 0.91 \pm 0.11(\text{exp.}) \pm 0.07(\text{theory})$
- Submitted to PRL, March 4th, 2009! <http://arxiv.org/abs/0903.0885>




CONCLUSIONS

- Single top searches are challenging
- Advance analysis tools essential to establish small signals buried underneath huge backgrounds
- **We report Observation (5σ) for electroweak single top quark at CDF**
 - $\sigma_{\text{SingleTop}} = 2.3^{+0.6}_{-0.5} \text{ pb}$
 - $|V_{tb}| = 0.91 \pm 0.11(\text{exp.}) \pm 0.07(\text{theory})$
- Future: properties in the single top sample
- Important milestone along the way to the Higgs!
 - All SM backgrounds for Higgs searches are now firmly established
- Check all the results in the public web page
http://www-cdf.fnal.gov/physics/new/top/public_singletop.html



BACKUP

SEARCH STRATEGY



Rate/shape systematic uncertainties: implemented as nuisance parameters

BACKGROUND ESTIMATE

Top/EWK (WW/WZ/Z $\rightarrow\tau\tau$, ttbar, single t)



- MC normalized to theoretical cross-section

W+HF jets (Wbb/Wcc/Wc)

- W+jets normalization from data and heavy flavor (HF) fractions from ALPGEN Monte Carlo, calibrated in generic multijet data


Non-W (QCD)

- Multijet events with semileptonic b -decays or mismeasured jets
- Fit low missing E_T data and extrapolate into signal region




Mistags (W+2jets)

- Falsely tagged light quark or gluon jets
- Mistag probability parameterization obtained from inclusive jet data




TEMPLATE SHAPE SYSTEMATICS




- A total of 370 shape uncertainties evaluated!
- Each template, each source of shape error, each channel (#tags, #jets, extended muon coverage)
- Shape uncertainties affect sensitivity - most are quite small but some appreciable

EXTENDED MUON COVERAGE



- Traditional muons are triggered muons with hits in
 - Both central muon chambers (CMUP, $|\eta| < 0.6$)
 - In the central extension chamber (CMX, $|\eta| < 1.0$)
- Extended muons
 - Events with missing $E_T > 35$ GeV plus two jets are triggered
 - Recover muons in the gaps left by the high p_T muon triggers
 - Acceptance gain of $\sim 30\%$
 - Orthogonal muon definitions

EXTENDED MUON COVERAGE




- Extended muons

- Events with missing $E_T > 35$ GeV plus two jets are triggered
- Recover muons in the gaps left by the high p_T muon triggers
- Acceptance gain of $\sim 30\%$
- Orthogonal muon definitions
 - CMU-only muons (η -Gap)
 - CMP-only muons (ϕ -Gap)
 - BMU muons
 - CMIO muons (stubless and no fiducial to any chamber)
 - SCMIO (stubbed and no match in other categories)
 - CMXNT (non-triggered CMX)


NN JET FLAVOR SEPARATOR

- About 50% of the background in the W+2 jets sample do NOT contain b quarks (even though a secondary vertex is required)
- Train Neural Network to separate b-quark jets from charm and light-quark jets
- Jet and track variables (vertex mass, decay length, track multiplicity...)
- Replace Yes-No tag decision by a continuous variable
- Improves single top sensitivity by $\sim 10\text{-}15\%$!




OVERVIEW OF BOOSTED DECISION TREES

- Decision Tree (DT): Sequence of binary splits using the discriminating variable which gives best sig-bkg separation
- Leaf nodes are classified as sig-like or bkg-like depending on majority of events ending up in the respective leaf
- A Boosting algorithm improves the discrimination power and statistical stability
 - Events misclassified during a DT training are given a higher weight in the next DT training
- To avoid overtraining a pruning algorithm (BDT) is performed after the BDT training
 - Removes statistically insignificant nodes by cutting back a tree from the bottom up



LIKELIHOOD FUNCTION OVERVIEW

- Combines several sensitive variables into a single one
- 7 (10) variables used in the 2 (3) jet bin: H_T , $Q \times \eta$, M_{jj} , $\cos(\theta_{l,j})$, $\log(ME_{t\text{-chan}})$...




$$p_i^{sig}(x_i) = \frac{N_i^{sig}(x_i)}{N_i^{sig}(x_i) + N_i^{bkg}(x_i)},$$

$$L = \frac{\prod_{i=1}^{nvar} p_i^{sig}(x_i)}{\prod_{i=1}^{nvar} p_i^{sig}(x_i) + \prod_{i=1}^{nvar} p_i^{bkg}(x_i)}$$

MATRIX ELEMENTS OVERVIEW

- Compute, for each event, the probability for signal and background hypotheses

- Use full event kinematic information
- Calculate probabilities for signal and backgrounds




- Build a discriminant

$$EPD = \frac{b \cdot P_{sig}(\vec{x})}{b \cdot P_{sig}(\vec{x}) + b \cdot P_{b-bkg}(\vec{x}) + (1-b) \cdot P_{nonb-bkg}(\vec{x})}$$


- b : b -likeness form NN Flavor Separator

PRL COMBINATION I




- 5 independent ℓ +jets analyses at CDF sharing single top selection (correlations 60–85%)
- Combined into a super discriminant trained using genetic algorithms to provide optimal sensitivity

PRL COMBINATION II




- Another orthogonal MET + jets analysis where the lepton is missed
- A simultaneous fit is performed to the super discriminant and the MET + jets channel
- Expected p-value: $> 5.9\sigma$
- **Observed p-value: 5.0σ**
- Measured cross-section: $\sigma = 2.3^{+0.6}_{-0.5}$ pb
- Extract $|V_{tb}| = 0.91 \pm 0.11(\text{exp.}) \pm 0.07(\text{the.})$
- Submitted to PRL, March 4th, 2009! <http://arxiv.org/abs/0903.0885>




DIRECT $|V_{tb}|$ MEASUREMENT

- Using cross section result measure $|V_{tb}|$
- Assume Standard Model (V-A) coupling and $|V_{tb}| \gg |V_{ts}|, |V_{td}|$
(from $\text{BR}(t \rightarrow Wb)$ measurements)



$$|V_{tb,meas}|^2 = \frac{\sigma_{meas}}{\sigma_{SM}} \cdot |V_{tb,SM}|^2$$






CDF combined fit:

$$|V_{tb}| > 0.71 \text{ at } 95\% \text{ C.L.}$$

Z. Sullivan, Phys.Rev. D70 (2004) 114012

$$|V_{tb}| = 0.91 \pm 0.11 \text{ (stat+syst)} \pm 0.07 \text{ (theory)}$$

SIGNAL FEATURES



- Super Discriminant > 0.72
- Purity $S/B \sim 1.2$