

Nonperturbative QFT

Romuald A. Janik

Jagiellonian University
Kraków

2009 Europhysics Conference on High Energy Physics
22 July 2009

- 1 Solving (nonperturbatively) QFT's**
 - The quintessential example: sine-Gordon
 - QCD???
 - $\mathcal{N} = 4$ Super-Yang-Mills theory
- 2 The AdS/CFT correspondence and $\mathcal{N} = 4$ SYM**
- 3 Why is it interesting?**
- 4 Methods at strong coupling**
- 5 Towards an exact solution for any coupling**
- 6 Example: Konishi anomalous dimension**
- 7 Outlook**

The quintessential example: sine-Gordon

- sine-Gordon theory is a **two-dimensional** QFT defined by the Lagrangian

$$S = \int d^2x \left\{ \frac{1}{2} \partial_\mu \Phi \partial^\mu \Phi + \frac{\mu^2}{\beta^2} \cos(\beta\Phi) \right\}$$

- Expanding $\cos(\beta\Phi)$ for small β generates an infinite set of vertices for Φ , one can develop perturbation theory...
- There exist solitons (and anti-solitons).. classical solutions, quadratic fluctuations...
- *But* the soliton-soliton S-matrix has been found analytically **for any** β

$$S(\theta) = - \exp \left\{ 2 \int_0^\infty \frac{dk}{k} \sin(k\theta) \frac{\sinh(p-1)k}{2 \cosh k \sinh pk} \right\}$$

where

$$p = \frac{\beta^2}{8\pi - \beta^2}$$

The quintessential example: sine-Gordon

- sine-Gordon theory is a **two-dimensional** QFT defined by the Lagrangian

$$S = \int d^2x \left\{ \frac{1}{2} \partial_\mu \Phi \partial^\mu \Phi + \frac{\mu^2}{\beta^2} \cos(\beta\Phi) \right\}$$

- Expanding $\cos(\beta\Phi)$ for small β generates an infinite set of vertices for Φ , one can develop perturbation theory...
- There exist solitons (and anti-solitons).. classical solutions, quadratic fluctuations...
- *But* the soliton-soliton S-matrix has been found analytically **for any** β

$$S(\theta) = - \exp \left\{ 2 \int_0^\infty \frac{dk}{k} \sin(k\theta) \frac{\sinh(p-1)k}{2 \cosh k \sinh pk} \right\}$$

where

$$p = \frac{\beta^2}{8\pi - \beta^2}$$

The quintessential example: sine-Gordon

- sine-Gordon theory is a **two-dimensional** QFT defined by the Lagrangian

$$S = \int d^2x \left\{ \frac{1}{2} \partial_\mu \Phi \partial^\mu \Phi + \frac{\mu^2}{\beta^2} \cos(\beta\Phi) \right\}$$

- Expanding $\cos(\beta\Phi)$ for small β generates an infinite set of vertices for Φ , one can develop perturbation theory...
- There exist solitons (and anti-solitons).. classical solutions, quadratic fluctuations...
- *But* the soliton-soliton S-matrix has been found analytically **for any** β

$$S(\theta) = - \exp \left\{ 2 \int_0^\infty \frac{dk}{k} \sin(k\theta) \frac{\sinh(p-1)k}{2 \cosh k \sinh pk} \right\}$$

where

$$p = \frac{\beta^2}{8\pi - \beta^2}$$

The quintessential example: sine-Gordon

- sine-Gordon theory is a **two-dimensional** QFT defined by the Lagrangian

$$S = \int d^2x \left\{ \frac{1}{2} \partial_\mu \Phi \partial^\mu \Phi + \frac{\mu^2}{\beta^2} \cos(\beta\Phi) \right\}$$

- Expanding $\cos(\beta\Phi)$ for small β generates an infinite set of vertices for Φ , one can develop perturbation theory...
- There exist solitons (and anti-solitons).. classical solutions, quadratic fluctuations...
- *But* the soliton-soliton S-matrix has been found analytically **for any** β

$$S(\theta) = - \exp \left\{ 2 \int_0^\infty \frac{dk}{k} \sin(k\theta) \frac{\sinh(p-1)k}{2 \cosh k \sinh pk} \right\}$$

where

$$p = \frac{\beta^2}{8\pi - \beta^2}$$

The quintessential example: sine-Gordon

- sine-Gordon theory is a **two-dimensional** QFT defined by the Lagrangian

$$S = \int d^2x \left\{ \frac{1}{2} \partial_\mu \Phi \partial^\mu \Phi + \frac{\mu^2}{\beta^2} \cos(\beta\Phi) \right\}$$

- Expanding $\cos(\beta\Phi)$ for small β generates an infinite set of vertices for Φ , one can develop perturbation theory...
- There exist solitons (and anti-solitons).. classical solutions, quadratic fluctuations...
- *But* the soliton-soliton S-matrix has been found analytically **for any** β

$$S(\theta) = - \exp \left\{ 2 \int_0^\infty \frac{dk}{k} \sin(k\theta) \frac{\sinh(p-1)k}{2 \cosh k \sinh pk} \right\}$$

where

$$p = \frac{\beta^2}{8\pi - \beta^2}$$

The quintessential example: sine-Gordon

- sine-Gordon theory is a **two-dimensional** QFT defined by the Lagrangian

$$S = \int d^2x \left\{ \frac{1}{2} \partial_\mu \Phi \partial^\mu \Phi + \frac{\mu^2}{\beta^2} \cos(\beta\Phi) \right\}$$

- Expanding $\cos(\beta\Phi)$ for small β generates an infinite set of vertices for Φ , one can develop perturbation theory...
- There exist solitons (and anti-solitons).. classical solutions, quadratic fluctuations...
- *But* the soliton-soliton S-matrix has been found analytically **for any** β

$$S(\theta) = - \exp \left\{ 2 \int_0^\infty \frac{dk}{k} \sin(k\theta) \frac{\sinh(p-1)k}{2 \cosh k \sinh pk} \right\}$$

where

$$p = \frac{\beta^2}{8\pi - \beta^2}$$

The quintessential example: sine-Gordon

- sine-Gordon theory is a **two-dimensional** QFT defined by the Lagrangian

$$S = \int d^2x \left\{ \frac{1}{2} \partial_\mu \Phi \partial^\mu \Phi + \frac{\mu^2}{\beta^2} \cos(\beta\Phi) \right\}$$

- Expanding $\cos(\beta\Phi)$ for small β generates an infinite set of vertices for Φ , one can develop perturbation theory...
- There exist solitons (and anti-solitons).. classical solutions, quadratic fluctuations...
- *But* the soliton-soliton S-matrix has been found analytically **for any** β

$$S(\theta) = - \exp \left\{ 2 \int_0^\infty \frac{dk}{k} \sin(k\theta) \frac{\sinh(p-1)k}{2 \cosh k \sinh pk} \right\}$$

where

$$p = \frac{\beta^2}{8\pi - \beta^2}$$

The quintessential example: sine-Gordon

- sine-Gordon theory is a **two-dimensional** QFT defined by the Lagrangian

$$S = \int d^2x \left\{ \frac{1}{2} \partial_\mu \Phi \partial^\mu \Phi + \frac{\mu^2}{\beta^2} \cos(\beta\Phi) \right\}$$

- Expanding $\cos(\beta\Phi)$ for small β generates an infinite set of vertices for Φ , one can develop perturbation theory...
- There exist solitons (and anti-solitons).. classical solutions, quadratic fluctuations...
- *But* the soliton-soliton S-matrix has been found analytically **for any** β

$$S(\theta) = - \exp \left\{ 2 \int_0^\infty \frac{dk}{k} \sin(k\theta) \frac{\sinh(p-1)k}{2 \cosh k \sinh pk} \right\}$$

where

$$p = \frac{\beta^2}{8\pi - \beta^2}$$

The quintessential example: sine-Gordon

- How is it possible??
- The sine-Gordon theory is an example of an **integrable quantum field theory**
- These theories possess additional higher spin/nonlocal conserved charges
- As a consequence the S-matrix factorizes into $2 \rightarrow 2$ scatterings and obeys Yang-Baxter Equation which allows to *nonperturbatively* find it exactly
- All this works **only** in two dimensions! No generalization to four-dimensional theories exists...

Two-dimensional integrable QFT's become relevant for four-dimensional physics due to the AdS/CFT correspondence

The quintessential example: sine-Gordon

- How is it possible??
- The sine-Gordon theory is an example of an **integrable quantum field theory**
- These theories possess additional higher spin/nonlocal conserved charges
- As a consequence the S-matrix factorizes into $2 \rightarrow 2$ scatterings and obeys Yang-Baxter Equation which allows to *nonperturbatively* find it exactly
- All this works **only** in two dimensions! No generalization to four-dimensional theories exists...

Two-dimensional integrable QFT's become relevant for four-dimensional physics due to the AdS/CFT correspondence

The quintessential example: sine-Gordon

- How is it possible??
- The sine-Gordon theory is an example of an **integrable quantum field theory**
- These theories possess additional higher spin/nonlocal conserved charges
- As a consequence the S-matrix factorizes into $2 \rightarrow 2$ scatterings and obeys Yang-Baxter Equation which allows to *nonperturbatively* find it exactly
- All this works **only** in two dimensions! No generalization to four-dimensional theories exists...

Two-dimensional integrable QFT's become relevant for four-dimensional physics due to the AdS/CFT correspondence

The quintessential example: sine-Gordon

- How is it possible??
- The sine-Gordon theory is an example of an **integrable quantum field theory**
- These theories possess additional higher spin/nonlocal conserved charges
- As a consequence the S-matrix factorizes into $2 \rightarrow 2$ scatterings and obeys Yang-Baxter Equation which allows to *nonperturbatively* find it exactly
- All this works **only** in two dimensions! No generalization to four-dimensional theories exists...

Two-dimensional integrable QFT's become relevant for four-dimensional physics due to the AdS/CFT correspondence

The quintessential example: sine-Gordon

- How is it possible??
- The sine-Gordon theory is an example of an **integrable quantum field theory**
- These theories possess additional higher spin/nonlocal conserved charges
- As a consequence the S-matrix factorizes into $2 \rightarrow 2$ scatterings and obeys Yang-Baxter Equation which allows to *nonperturbatively* find it exactly
- All this works **only** in two dimensions! No generalization to four-dimensional theories exists...

Two-dimensional integrable QFT's become relevant for four-dimensional physics due to the AdS/CFT correspondence

The quintessential example: sine-Gordon

- How is it possible??
- The sine-Gordon theory is an example of an **integrable quantum field theory**
- These theories possess additional higher spin/nonlocal conserved charges
- As a consequence the S-matrix factorizes into $2 \rightarrow 2$ scatterings and obeys Yang-Baxter Equation which allows to *nonperturbatively* find it exactly
- All this works **only** in two dimensions! No generalization to four-dimensional theories exists...

Two-dimensional integrable QFT's become relevant for four-dimensional physics due to the AdS/CFT correspondence

The quintessential example: sine-Gordon

- How is it possible??
- The sine-Gordon theory is an example of an **integrable quantum field theory**
- These theories possess additional higher spin/nonlocal conserved charges
- As a consequence the S-matrix factorizes into $2 \rightarrow 2$ scatterings and obeys Yang-Baxter Equation which allows to *nonperturbatively* find it exactly
- All this works **only** in two dimensions! No generalization to four-dimensional theories exists...

Two-dimensional integrable QFT's become relevant for **four-dimensional** physics due to the AdS/CFT correspondence

QCD

- Inherently difficult...
- Running coupling
- Mixture of perturbative and nonperturbative effects...

Traditional methods:

- Lattice QCD (inherently Euclidean, 'black box')
- Models of QCD vacuum
- Effective models (not *derivable* from the fundamental theory)

Look at other, 'simpler' gauge theories for which other methods exist...

QCD

- Inherently difficult...
- Running coupling
- Mixture of perturbative and nonperturbative effects...

Traditional methods:

- Lattice QCD (inherently Euclidean, 'black box')
- Models of QCD vacuum
- Effective models (not *derivable* from the fundamental theory)

Look at other, 'simpler' gauge theories for which other methods exist...

QCD

- Inherently difficult...
- Running coupling
- Mixture of perturbative and nonperturbative effects...

Traditional methods:

- Lattice QCD (inherently Euclidean, 'black box')
- Models of QCD vacuum
- Effective models (not *derivable* from the fundamental theory)

Look at other, 'simpler' gauge theories for which other methods exist...

QCD

- Inherently difficult...
- Running coupling
- Mixture of perturbative and nonperturbative effects...

Traditional methods:

- Lattice QCD (inherently Euclidean, 'black box')
- Models of QCD vacuum
- Effective models (not *derivable* from the fundamental theory)

Look at other, 'simpler' gauge theories for which other methods exist...

QCD

- Inherently difficult...
- Running coupling
- Mixture of perturbative and nonperturbative effects...

Traditional methods:

- Lattice QCD (inherently Euclidean, 'black box')
- Models of QCD vacuum
- Effective models (not *derivable* from the fundamental theory)

Look at other, 'simpler' gauge theories for which other methods exist...

QCD

- Inherently difficult...
- Running coupling
- Mixture of perturbative and nonperturbative effects...

Traditional methods:

- Lattice QCD (inherently Euclidean, 'black box')
- Models of QCD vacuum
- Effective models (not *derivable* from the fundamental theory)

Look at other, 'simpler' gauge theories for which other methods exist...

QCD

- Inherently difficult...
- Running coupling
- Mixture of perturbative and nonperturbative effects...

Traditional methods:

- Lattice QCD (inherently Euclidean, 'black box')
- Models of QCD vacuum
- Effective models (not *derivable* from the fundamental theory)

Look at other, 'simpler' gauge theories for which other methods exist...

QCD

- Inherently difficult...
- Running coupling
- Mixture of perturbative and nonperturbative effects...

Traditional methods:

- Lattice QCD (inherently Euclidean, 'black box')
- Models of QCD vacuum
- Effective models (not *derivable* from the fundamental theory)

Look at other, 'simpler' gauge theories for which other methods exist...

QCD

- Inherently difficult...
- Running coupling
- Mixture of perturbative and nonperturbative effects...

Traditional methods:

- Lattice QCD (inherently Euclidean, 'black box')
- Models of QCD vacuum
- Effective models (not *derivable* from the fundamental theory)

Look at other, 'simpler' gauge theories for which other methods exist...

QCD

- Inherently difficult...
- Running coupling
- Mixture of perturbative and nonperturbative effects...

Traditional methods:

- Lattice QCD (inherently Euclidean, 'black box')
- Models of QCD vacuum
- Effective models (*not derivable* from the fundamental theory)

Look at other, 'simpler' gauge theories for which other methods exist...

QCD

- Inherently difficult...
- Running coupling
- Mixture of perturbative and nonperturbative effects...

Traditional methods:

- Lattice QCD (inherently Euclidean, 'black box')
- Models of QCD vacuum
- Effective models (not *derivable* from the fundamental theory)

Look at other, 'simpler' gauge theories for which other methods exist...

QCD

- Inherently difficult...
- Running coupling
- Mixture of perturbative and nonperturbative effects...

Traditional methods:

- Lattice QCD (inherently Euclidean, 'black box')
- Models of QCD vacuum
- Effective models (not *derivable* from the fundamental theory)

Look at other, 'simpler' gauge theories for which other methods exist...

Lot of progress has been made in the understanding of $\mathcal{N} = 4$ Super-Yang-Mills theory...

- $\mathcal{N} = 4$ Super Yang-Mills (\equiv ordinary Yang-Mills+4 adjoint fermions +6 adjoint scalars+ appropriate self-interactions)
- This theory is
 - 1 supersymmetric
 - 2 conformal (scale-invariant even at the quantum level)
- **Bonus:** There exists an efficient way of studying this theory at strong coupling: the AdS/CFT (Anti-deSitter/Conformal Field Theory) correspondence

Lot of progress has been made in the understanding of $\mathcal{N} = 4$ Super-Yang-Mills theory...

- $\mathcal{N} = 4$ Super Yang-Mills (\equiv ordinary Yang-Mills+4 adjoint fermions +6 adjoint scalars+ appropriate self-interactions)
- This theory is
 - 1 supersymmetric
 - 2 conformal (scale-invariant even at the quantum level)
- **Bonus:** There exists an efficient way of studying this theory at strong coupling: the AdS/CFT (Anti-deSitter/Conformal Field Theory) correspondence

Lot of progress has been made in the understanding of $\mathcal{N} = 4$ Super-Yang-Mills theory...

- $\mathcal{N} = 4$ Super Yang-Mills (\equiv ordinary Yang-Mills+4 adjoint fermions +6 adjoint scalars+ appropriate self-interactions)
- This theory is
 - 1 supersymmetric
 - 2 conformal (scale-invariant even at the quantum level)
- **Bonus:** There exists an efficient way of studying this theory at strong coupling: the AdS/CFT (Anti-deSitter/Conformal Field Theory) correspondence

Lot of progress has been made in the understanding of $\mathcal{N} = 4$ Super-Yang-Mills theory...

- $\mathcal{N} = 4$ Super Yang-Mills (\equiv ordinary Yang-Mills+4 adjoint fermions +6 adjoint scalars+ appropriate self-interactions)
- This theory is
 - 1 supersymmetric
 - 2 conformal (scale-invariant even at the quantum level)
- **Bonus:** There exists an efficient way of studying this theory at strong coupling: the AdS/CFT (Anti-deSitter/Conformal Field Theory) correspondence

Lot of progress has been made in the understanding of $\mathcal{N} = 4$ Super-Yang-Mills theory...

- $\mathcal{N} = 4$ Super Yang-Mills (\equiv ordinary Yang-Mills+4 adjoint fermions +6 adjoint scalars+ appropriate self-interactions)
- This theory is
 - 1 supersymmetric
 - 2 conformal (scale-invariant even at the quantum level)
- **Bonus:** There exists an efficient way of studying this theory at strong coupling: the AdS/CFT (Anti-deSitter/Conformal Field Theory) correspondence

Lot of progress has been made in the understanding of $\mathcal{N} = 4$ Super-Yang-Mills theory...

- $\mathcal{N} = 4$ Super Yang-Mills (\equiv ordinary Yang-Mills+4 adjoint fermions +6 adjoint scalars+ appropriate self-interactions)
- This theory is
 - 1 supersymmetric
 - 2 conformal (scale-invariant even at the quantum level)
- **Bonus:** There exists an efficient way of studying this theory at strong coupling: the AdS/CFT (Anti-deSitter/Conformal Field Theory) correspondence

Lot of progress has been made in the understanding of $\mathcal{N} = 4$ Super-Yang-Mills theory...

- $\mathcal{N} = 4$ Super Yang-Mills (\equiv ordinary Yang-Mills+4 adjoint fermions +6 adjoint scalars+ appropriate self-interactions)
- This theory is
 - 1 supersymmetric
 - 2 conformal (scale-invariant even at the quantum level)
- **Bonus:** There exists an efficient way of studying this theory at strong coupling: the AdS/CFT (Anti-deSitter/Conformal Field Theory) correspondence

Lot of progress has been made in the understanding of $\mathcal{N} = 4$ Super-Yang-Mills theory...

- $\mathcal{N} = 4$ Super Yang-Mills (\equiv ordinary Yang-Mills+4 adjoint fermions +6 adjoint scalars+ appropriate self-interactions)
- This theory is
 - 1 supersymmetric
 - 2 conformal (scale-invariant even at the quantum level)
- **Bonus:** There exists an efficient way of studying this theory at strong coupling: the AdS/CFT (Anti-deSitter/Conformal Field Theory) correspondence

Lot of progress has been made in the understanding of $\mathcal{N} = 4$ Super-Yang-Mills theory...

- $\mathcal{N} = 4$ Super Yang-Mills (\equiv ordinary Yang-Mills+4 adjoint fermions +6 adjoint scalars+ appropriate self-interactions)
- This theory is
 - 1 supersymmetric
 - 2 conformal (scale-invariant even at the quantum level)
- **Bonus:** There exists an efficient way of studying this theory at strong coupling: the AdS/CFT (Anti-deSitter/Conformal Field Theory) correspondence

$\mathcal{N} = 4$ Super Yang-Mills theory

\equiv

Superstrings on $AdS_5 \times S^5$

strong coupling
nonperturbative physics

very difficult

weak coupling
'easy'

(semi-)classical strings
or supergravity

'easy'

highly quantum regime
very difficult

- New ways of looking at nonperturbative gauge theory physics...
- Intricate links with General Relativity...

$\mathcal{N} = 4$ Super Yang-Mills theory

\equiv

Superstrings on $AdS_5 \times S^5$

strong coupling
nonperturbative physics

very difficult

weak coupling
'easy'

(semi-)classical strings
or supergravity

'easy'

highly quantum regime
very difficult

- New ways of looking at nonperturbative gauge theory physics...
- Intricate links with General Relativity...

$\mathcal{N} = 4$ Super Yang-Mills theory

\equiv

Superstrings on $AdS_5 \times S^5$

strong coupling
nonperturbative physics

very difficult

weak coupling
'easy'

(semi-)classical strings
or supergravity

'easy'

highly quantum regime
very difficult

- New ways of looking at nonperturbative gauge theory physics...
- Intricate links with General Relativity...

More theoretical perspective:

- Fascinating as it relates two *completely different* theoretical constructions: 4D gauge theory and string theory in 10D
- Use $\mathcal{N} = 4$ SYM as a theoretical laboratory for studying nonperturbative gauge theory physics
- In this theory one can perform quite rigorous computations at strong coupling
- The natural language of the AdS/CFT correspondence appropriate to strongly coupled $\mathcal{N} = 4$ SYM is quite new w.r.t. conventional gauge theory methods
- Try to build some new physical intuitions within this new language

$\mathcal{N} = 4$ SYM may be the 'harmonic oscillator' of four dimensional gauge theories
D. Gross

Why is it interesting?

More theoretical perspective:

- Fascinating as it relates two *completely different* theoretical constructions: 4D gauge theory and string theory in 10D
- Use $\mathcal{N} = 4$ SYM as a theoretical laboratory for studying nonperturbative gauge theory physics
- In this theory one can perform quite rigorous computations at strong coupling
- The natural language of the AdS/CFT correspondence appropriate to strongly coupled $\mathcal{N} = 4$ SYM is quite new w.r.t. conventional gauge theory methods
- Try to build some new physical intuitions within this new language

$\mathcal{N} = 4$ SYM may be the 'harmonic oscillator' of four dimensional gauge theories

D. Gross

Why is it interesting?

More theoretical perspective:

- Fascinating as it relates two *completely different* theoretical constructions: 4D gauge theory and string theory in 10D
- Use $\mathcal{N} = 4$ SYM as a theoretical laboratory for studying nonperturbative gauge theory physics
- In this theory one can perform quite rigorous computations at strong coupling
- The natural language of the AdS/CFT correspondence appropriate to strongly coupled $\mathcal{N} = 4$ SYM is quite new w.r.t. conventional gauge theory methods
- Try to build some new physical intuitions within this new language

$\mathcal{N} = 4$ SYM may be the 'harmonic oscillator' of four dimensional gauge theories

D. Gross

Why is it interesting?

More theoretical perspective:

- Fascinating as it relates two *completely different* theoretical constructions: 4D gauge theory and string theory in 10D
- Use $\mathcal{N} = 4$ SYM as a theoretical laboratory for studying nonperturbative gauge theory physics
- In this theory one can perform quite rigorous computations at strong coupling
- The natural language of the AdS/CFT correspondence appropriate to strongly coupled $\mathcal{N} = 4$ SYM is quite new w.r.t. conventional gauge theory methods
- Try to build some new physical intuitions within this new language

$\mathcal{N} = 4$ SYM may be the 'harmonic oscillator' of four dimensional gauge theories

D. Gross

Why is it interesting?

More theoretical perspective:

- Fascinating as it relates two *completely different* theoretical constructions: 4D gauge theory and string theory in 10D
- Use $\mathcal{N} = 4$ SYM as a theoretical laboratory for studying nonperturbative gauge theory physics
- In this theory one can perform quite rigorous computations at strong coupling
- The natural language of the AdS/CFT correspondence appropriate to strongly coupled $\mathcal{N} = 4$ SYM is quite new w.r.t. conventional gauge theory methods
- Try to build some new physical intuitions within this new language

$\mathcal{N} = 4$ SYM may be the 'harmonic oscillator' of four dimensional gauge theories

D. Gross

More theoretical perspective:

- Fascinating as it relates two *completely different* theoretical constructions: 4D gauge theory and string theory in 10D
- Use $\mathcal{N} = 4$ SYM as a theoretical laboratory for studying nonperturbative gauge theory physics
- In this theory one can perform quite rigorous computations at strong coupling
- The natural language of the AdS/CFT correspondence appropriate to strongly coupled $\mathcal{N} = 4$ SYM is quite new w.r.t. conventional gauge theory methods
- Try to build some new physical intuitions within this new language

$\mathcal{N} = 4$ SYM may be the 'harmonic oscillator' of four dimensional gauge theories

D. Gross

Why is it interesting?

More theoretical perspective:

- Fascinating as it relates two *completely different* theoretical constructions: 4D gauge theory and string theory in 10D
- Use $\mathcal{N} = 4$ SYM as a theoretical laboratory for studying nonperturbative gauge theory physics
- In this theory one can perform quite rigorous computations at strong coupling
- The natural language of the AdS/CFT correspondence appropriate to strongly coupled $\mathcal{N} = 4$ SYM is quite new w.r.t. conventional gauge theory methods
- Try to build some new physical intuitions within this new language

$\mathcal{N} = 4$ SYM may be the 'harmonic oscillator' of four dimensional gauge theories

D. Gross

More practical perspective:

- Although its behaviour is quite different from QCD at $T = 0$ (no confinement etc.), at *nonzero* temperature supersymmetry is automatically broken, the theory is deconfined
- Use the results on strong coupling properties of $\mathcal{N} = 4$ plasma as a point of reference for analyzing/describing QCD plasma
 - see talk by R. Peschanski in the Heavy-Ion session
 - see plenary talk by U. Wiedemann later today
- Works directly in Minkowski signature
- In particular many gauge-theoretical problems are translated into geometrical General Relativity like questions which are tractable (even analytically)

More practical perspective:

- Although its behaviour is quite different from QCD at $T = 0$ (no confinement etc.), at *nonzero* temperature supersymmetry is automatically broken, the theory is deconfined
- Use the results on strong coupling properties of $\mathcal{N} = 4$ plasma as a point of reference for analyzing/describing QCD plasma
 - see talk by R. Peschanski in the Heavy-Ion session
 - see plenary talk by U. Wiedemann later today
- Works directly in Minkowski signature
- In particular many gauge-theoretical problems are translated into geometrical General Relativity like questions which are tractable (even analytically)

More practical perspective:

- Although its behaviour is quite different from QCD at $T = 0$ (no confinement etc.), at *nonzero* temperature supersymmetry is automatically broken, the theory is deconfined
- Use the results on strong coupling properties of $\mathcal{N} = 4$ plasma as a point of reference for analyzing/describing QCD plasma
 - see talk by R. Peschanski in the Heavy-Ion session
 - see plenary talk by U. Wiedemann later today
- Works directly in Minkowski signature
- In particular many gauge-theoretical problems are translated into geometrical General Relativity like questions which are tractable (even analytically)

More practical perspective:

- Although its behaviour is quite different from QCD at $T = 0$ (no confinement etc.), at *nonzero* temperature supersymmetry is automatically broken, the theory is deconfined
- Use the results on strong coupling properties of $\mathcal{N} = 4$ plasma as a point of reference for analyzing/describing QCD plasma
 - see talk by R. Peschanski in the Heavy-Ion session
 - see plenary talk by U. Wiedemann later today
- Works directly in Minkowski signature
- In particular many gauge-theoretical problems are translated into geometrical General Relativity like questions which are tractable (even analytically)

More practical perspective:

- Although its behaviour is quite different from QCD at $T = 0$ (no confinement etc.), at *nonzero* temperature supersymmetry is automatically broken, the theory is deconfined
- Use the results on strong coupling properties of $\mathcal{N} = 4$ plasma as a point of reference for analyzing/describing QCD plasma
 - see talk by R. Peschanski in the Heavy-Ion session
 - see plenary talk by U. Wiedemann later today
- Works directly in Minkowski signature
- In particular many gauge-theoretical problems are translated into geometrical General Relativity like questions which are tractable (even analytically)

What degrees of freedom appear on the string side???

- Vibrational modes of the string \equiv particles/fields in $AdS_5 \times S^5$
- Massless modes \equiv graviton+...
- At **strong** coupling massive modes are very heavy \rightarrow it is enough to restrict oneself to dynamics of massless modes \equiv gravity
- At weaker coupling massive string modes become important...

Is there hope for an exact solution at any coupling???

What degrees of freedom appear on the string side???

- Vibrational modes of the string \equiv particles/fields in $AdS_5 \times S^5$
- Massless modes \equiv graviton+...
- At **strong** coupling massive modes are very heavy \rightarrow it is enough to restrict oneself to dynamics of massless modes \equiv gravity
- At weaker coupling massive string modes become important...

Is there hope for an exact solution at any coupling???

What degrees of freedom appear on the string side???

- Vibrational modes of the string \equiv particles/fields in $AdS_5 \times S^5$
- Massless modes \equiv graviton+...
- At **strong** coupling massive modes are very heavy \rightarrow it is enough to restrict oneself to dynamics of massless modes \equiv gravity
- At weaker coupling massive string modes become important...

Is there hope for an exact solution at any coupling???

What degrees of freedom appear on the string side???

- Vibrational modes of the string \equiv particles/fields in $AdS_5 \times S^5$
- Massless modes \equiv graviton+...
- At **strong** coupling massive modes are very heavy \longrightarrow it is enough to restrict oneself to dynamics of massless modes \equiv gravity
- At weaker coupling massive string modes become important...

Is there hope for an exact solution at any coupling???

What degrees of freedom appear on the string side???

- Vibrational modes of the string \equiv particles/fields in $AdS_5 \times S^5$
- Massless modes \equiv graviton+...
- At **strong** coupling massive modes are very heavy \rightarrow it is enough to restrict oneself to dynamics of massless modes \equiv gravity
- At weaker coupling massive string modes become important...

Is there hope for an exact solution at any coupling???

What degrees of freedom appear on the string side???

- Vibrational modes of the string \equiv particles/fields in $AdS_5 \times S^5$
- Massless modes \equiv graviton+...
- At **strong** coupling massive modes are very heavy \rightarrow it is enough to restrict oneself to dynamics of massless modes \equiv gravity
- At weaker coupling massive string modes become important...

Is there hope for an exact solution at any coupling???

What degrees of freedom appear on the string side???

- Vibrational modes of the string \equiv particles/fields in $AdS_5 \times S^5$
- Massless modes \equiv graviton+...
- At **strong** coupling massive modes are very heavy \rightarrow it is enough to restrict oneself to dynamics of massless modes \equiv gravity
- At weaker coupling massive string modes become important...

Is there hope for an exact solution at any coupling???

How to describe strings in $AdS_5 \times S^5$?

- Consider a closed string in $AdS_5 \times S^5$:
- The embedding coordinates of the point (τ, σ) are *quantum fields* $X^\mu(\tau, \sigma)$ on the worldsheet which has the geometry of a cylinder
- String theory in $AdS_5 \times S^5 \equiv$ a specific two dimensional quantum field theory defined on a cylinder (worldsheet QFT)
- It turns out that this worldsheet QFT is *integrable* and one can expect to solve this theory *exactly* for any coupling! (recall sine-Gordon...)

How to describe strings in $AdS_5 \times S^5$?

- Consider a closed string in $AdS_5 \times S^5$:
- The embedding coordinates of the point (τ, σ) are *quantum fields* $X^\mu(\tau, \sigma)$ on the worldsheet which has the geometry of a cylinder
- String theory in $AdS_5 \times S^5 \equiv$ a specific two dimensional quantum field theory defined on a cylinder (worldsheet QFT)
- It turns out that this worldsheet QFT is *integrable* and one can expect to solve this theory *exactly* for any coupling! (recall sine-Gordon...)

How to describe strings in $AdS_5 \times S^5$?

- Consider a closed string in $AdS_5 \times S^5$:



- The embedding coordinates of the point (τ, σ) are *quantum fields* $X^\mu(\tau, \sigma)$ on the worldsheet which has the geometry of a cylinder
- String theory in $AdS_5 \times S^5 \equiv$ a specific two dimensional quantum field theory defined on a cylinder (worldsheet QFT)
- It turns out that this worldsheet QFT is *integrable* and one can expect to solve this theory *exactly* for any coupling! (recall sine-Gordon...)

How to describe strings in $AdS_5 \times S^5$?

- Consider a closed string in $AdS_5 \times S^5$:



- The embedding coordinates of the point (τ, σ) are *quantum fields* $X^\mu(\tau, \sigma)$ on the worldsheet which has the geometry of a cylinder
- String theory in $AdS_5 \times S^5 \equiv$ a specific two dimensional quantum field theory defined on a cylinder (worldsheet QFT)
- It turns out that this worldsheet QFT is *integrable* and one can expect to solve this theory *exactly* for any coupling! (recall sine-Gordon...)

How to describe strings in $AdS_5 \times S^5$?

- Consider a closed string in $AdS_5 \times S^5$:



- The embedding coordinates of the point (τ, σ) are *quantum fields* $X^\mu(\tau, \sigma)$ on the worldsheet which has the geometry of a cylinder
- String theory in $AdS_5 \times S^5 \equiv$ a specific two dimensional quantum field theory defined on a cylinder (worldsheet QFT)
- It turns out that this worldsheet QFT is *integrable* and one can expect to solve this theory *exactly* for any coupling! (recall sine-Gordon...)

How to describe strings in $AdS_5 \times S^5$?

- Consider a closed string in $AdS_5 \times S^5$:



- The embedding coordinates of the point (τ, σ) are *quantum fields* $X^\mu(\tau, \sigma)$ on the worldsheet which has the geometry of a cylinder
- String theory in $AdS_5 \times S^5 \equiv$ a specific two dimensional quantum field theory defined on a cylinder (worldsheet QFT)
- It turns out that this worldsheet QFT is *integrable* and one can expect to solve this theory *exactly* for any coupling! (recall sine-Gordon...)

Towards an exact solution for any coupling

What do we mean by 'solve'?

Local operators
in gauge theory



String states
in $AdS_5 \times S^5$

Operator dimension

$$\langle O(x)O(y) \rangle = \frac{const}{|x-y|^{2\Delta}}$$



Energy of the corresponding
string state

One has to find the energy levels of an *integrable* two-dimensional QFT...

What do we mean by 'solve'?

Local operators
in gauge theory



String states
in $AdS_5 \times S^5$

Operator dimension

$$\langle O(x)O(y) \rangle = \frac{const}{|x - y|^{2\Delta}}$$



Energy of the corresponding
string state

One has to find the energy levels of an *integrable* two-dimensional QFT...

What do we mean by 'solve'?

Local operators
in gauge theory



String states
in $AdS_5 \times S^5$

Operator dimension

$$\langle O(x)O(y) \rangle = \frac{const}{|x - y|^{2\Delta}}$$



Energy of the corresponding
string state

One has to find the energy levels of an *integrable* two-dimensional QFT...

What do we mean by 'solve'?

Local operators
in gauge theory



String states
in $AdS_5 \times S^5$

Operator dimension

$$\langle O(x)O(y) \rangle = \frac{const}{|x - y|^{2\Delta}}$$



Energy of the corresponding
string state

One has to find the energy levels of an *integrable* two-dimensional QFT...

Example: Konishi anomalous dimension

- Simplest nonprotected operator: the Konishi operator

$$\text{tr } \Phi_j^2 \longleftrightarrow \text{tr } Z^2 X^2 + \dots \longleftrightarrow \text{tr } Z D^2 Z + \dots$$

- When computing anomalous dimensions from two point functions there are two types of graphs:
and

- The first class is contained in the so-called Asymptotic Bethe Ansatz of Beisert and Staudacher
- The second class are 'wrapping interactions' which start to appear at order g^{2L} (these are **not** contained in the Asymptotic Bethe Ansatz)
- Last year these wrapping graphs were computed at 4 loops by F.Fiamberti, A.Santambrogio, C.Sieg and D.Zanon

Example: Konishi anomalous dimension

- Simplest nonprotected operator: the Konishi operator

$$\text{tr } \Phi_i^2 \longleftrightarrow \text{tr } Z^2 X^2 + \dots \longleftrightarrow \text{tr } Z D^2 Z + \dots$$

- When computing anomalous dimensions from two point functions there are two types of graphs:

and


- The first class is contained in the so-called Asymptotic Bethe Ansatz of Beisert and Staudacher
- The second class are 'wrapping interactions' which start to appear at order g^{2L} (these are **not** contained in the Asymptotic Bethe Ansatz)
- Last year these wrapping graphs were computed at 4 loops by F.Fiamberti, A.Santambrogio, C.Sieg and D.Zanon

Example: Konishi anomalous dimension


- Simplest nonprotected operator: the Konishi operator

$$\text{tr } \Phi_i^2 \longleftrightarrow \text{tr } Z^2 X^2 + \dots \longleftrightarrow \text{tr } Z D^2 Z + \dots$$

- When computing anomalous dimensions from two point functions there are two types of graphs:



and




- The first class is contained in the so-called Asymptotic Bethe Ansatz of Beisert and Staudacher
- The second class are 'wrapping interactions' which start to appear at order g^{2L} (these are **not** contained in the Asymptotic Bethe Ansatz)
- Last year these wrapping graphs were computed at 4 loops by F.Fiamberti, A.Santambrogio, C.Sieg and D.Zanon

Example: Konishi anomalous dimension


- Simplest nonprotected operator: the Konishi operator

$$\text{tr } \Phi_i^2 \longleftrightarrow \text{tr } Z^2 X^2 + \dots \longleftrightarrow \text{tr } Z D^2 Z + \dots$$

- When computing anomalous dimensions from two point functions there are two types of graphs:



and




- The first class is contained in the so-called Asymptotic Bethe Ansatz of Beisert and Staudacher
- The second class are 'wrapping interactions' which start to appear at order g^{2L} (these are **not** contained in the Asymptotic Bethe Ansatz)
- Last year these wrapping graphs were computed at 4 loops by F.Fiamberti, A.Santambrogio, C.Sieg and D.Zanon

Example: Konishi anomalous dimension


- Simplest nonprotected operator: the Konishi operator

$$\text{tr } \Phi_i^2 \longleftrightarrow \text{tr } Z^2 X^2 + \dots \longleftrightarrow \text{tr } Z D^2 Z + \dots$$

- When computing anomalous dimensions from two point functions there are two types of graphs:



and




- The first class is contained in the so-called Asymptotic Bethe Ansatz of Beisert and Staudacher
- The second class are '**wrapping interactions**' which start to appear at order g^{2L} (these are **not** contained in the Asymptotic Bethe Ansatz)
- Last year these wrapping graphs were computed at 4 loops by F.Fiamberti, A.Santambrogio, C.Sieg and D.Zanon

Example: Konishi anomalous dimension


- Simplest nonprotected operator: the Konishi operator

$$\text{tr } \Phi_i^2 \longleftrightarrow \text{tr } Z^2 X^2 + \dots \longleftrightarrow \text{tr } Z D^2 Z + \dots$$

- When computing anomalous dimensions from two point functions there are two types of graphs:



and




- The first class is contained in the so-called Asymptotic Bethe Ansatz of Beisert and Staudacher
- The second class are '**wrapping interactions**' which start to appear at order g^{2L} (these are **not** contained in the Asymptotic Bethe Ansatz)
- Last year these wrapping graphs were computed at 4 loops by F.Fiamberti, A.Santambrogio, C.Sieg and D.Zanon

Example: Konishi anomalous dimension


- Simplest nonprotected operator: the Konishi operator

$$\text{tr } \Phi_i^2 \longleftrightarrow \text{tr } Z^2 X^2 + \dots \longleftrightarrow \text{tr } Z D^2 Z + \dots$$

- When computing anomalous dimensions from two point functions there are two types of graphs:



and



- The first class is contained in the so-called Asymptotic Bethe Ansatz of Beisert and Staudacher
- The second class are 'wrapping interactions' which start to appear at order g^{2L} (these are **not** contained in the Asymptotic Bethe Ansatz)
- Last year these wrapping graphs were computed at 4 loops by **F.Fiamberti**, **A.Santambrogio**, **C.Sieg** and **D.Zanon**




Figure C.1: Wrapping diagrams with chiral structure $\chi(1, 2, 3)$




Figure C.2: Wrapping diagrams with chiral structure $\chi(1, 3, 2)$

$W_{C1} \rightarrow *$	1	$W_{C4} \rightarrow \text{finite}$	
$W_{C2} \rightarrow *$	2	$W_{C5} \rightarrow -W_{C3}$	
$W_{C3} \rightarrow -W_{C5}$		$W_{C6} \rightarrow \text{finite}$	

Table C.2: Results of D -algebra for diagrams with structure $\chi(1, 3, 2)$







Figure C.3: Wrapping diagrams with chiral structure $\chi(2, 1, 3)$










Figure C.6: Wrapping diagrams with chiral structure $\chi(1)$ (continued)

$$\begin{aligned}
I_1 = J_1 &= \text{Diagram 1} = \frac{1}{(4\pi)^8} \left(-\frac{1}{24\varepsilon^4} + \frac{1}{4\varepsilon^3} - \frac{19}{24\varepsilon^2} + \frac{5}{4\varepsilon} \right) \\
I_2 &= \text{Diagram 2} = \frac{1}{(4\pi)^8} \left(-\frac{1}{24\varepsilon^4} + \frac{1}{4\varepsilon^3} - \frac{19}{24\varepsilon^2} + \frac{1}{\varepsilon} \left(\frac{5}{4} - \zeta(3) \right) \right) \\
I_3 = J_5 &= \text{Diagram 3} = \frac{1}{(4\pi)^8} \left(-\frac{1}{12\varepsilon^4} + \frac{1}{3\varepsilon^3} - \frac{5}{12\varepsilon^2} - \frac{1}{\varepsilon} \left(\frac{1}{2} - \zeta(3) \right) \right) \\
I_4 &= \text{Diagram 4} = \frac{1}{(4\pi)^8} \left(-\frac{1}{6\varepsilon^4} + \frac{1}{3\varepsilon^3} + \frac{1}{3\varepsilon^2} - \frac{1}{\varepsilon} (1 - \zeta(3)) \right) \\
I_5 &= \text{Diagram 5} = \frac{1}{(4\pi)^8} \frac{1}{\varepsilon} 5\zeta(5) \\
I_6 &= \text{Diagram 6} = \frac{1}{(4\pi)^8} \left(\frac{1}{12\varepsilon^2} - \frac{7}{12\varepsilon} \right) \quad I_7 = \text{Diagram 7} = \frac{1}{(4\pi)^8} \frac{1}{\varepsilon} (-\zeta(3)) \\
I_8 &= \text{Diagram 8} = \frac{1}{(4\pi)^8} \left(\frac{1}{4\varepsilon^2} - \frac{11}{12\varepsilon} \right) \quad I_9 = \text{Diagram 9} = \frac{1}{(4\pi)^8} \frac{1}{\varepsilon} \left(\frac{1}{2} \zeta(3) - \frac{5}{2} \zeta(5) \right) \\
I_{10} &= \text{Diagram 10} = \frac{1}{(4\pi)^8} \frac{1}{\varepsilon} \left(-\frac{1}{2} - \frac{1}{2} \zeta(3) + \frac{5}{2} \zeta(5) \right) \\
I_{11} &= \text{Diagram 11} = \frac{1}{(4\pi)^8} \frac{1}{\varepsilon} \left(-\frac{1}{4} - \frac{3}{2} \zeta(3) + \frac{5}{2} \zeta(5) \right) \\
I_{12} &= \text{Diagram 12} = \frac{1}{(4\pi)^8} \frac{1}{\varepsilon} \left(-\frac{1}{8} - \frac{1}{4} \zeta(3) + \frac{5}{4} \zeta(5) \right)
\end{aligned}$$

Table C.8: Loop integrals for 4-loop wrapping diagrams. The arrows of the same type indicate contracted spacetime derivatives

- The final result for the wrapping part at 4 loops is

$$\Delta_{wrapping} E = (324 + 864\zeta(3) - 1440\zeta(5))g^8$$

Compute the same 4-loop
anomalous dimension from
string theory

- The final result for the wrapping part at 4 loops is

$$\Delta_{wrapping} E = (324 + 864\zeta(3) - 1440\zeta(5))g^8$$

Compute the same 4-loop
anomalous dimension from
string theory

- The Konishi operator corresponds to a two particle state in the 2D worldsheet QFT of the string in $AdS_5 \times S^5$
- The wrapping graphs contributing at 4-loops correspond to a *single* 'virtual' graph:

- This corresponds to a simple expression

$$\Delta_w^{(4-loop)} = - \sum_{Q=1}^{\infty} \int_{-\infty}^{\infty} \frac{dq}{2\pi} Y_Q(q)$$

where $Y_Q(q)$ is a relatively simple rational function of q and Q


- The Konishi operator corresponds to a two particle state in the 2D worldsheet QFT of the string in $AdS_5 \times S^5$
- The wrapping graphs contributing at 4-loops correspond to a *single* 'virtual' graph:

- This corresponds to a simple expression

$$\Delta_w^{(4-loop)} = - \sum_{Q=1}^{\infty} \int_{-\infty}^{\infty} \frac{dq}{2\pi} Y_Q(q)$$

where $Y_Q(q)$ is a relatively simple rational function of q and Q

- The Konishi operator corresponds to a two particle state in the 2D worldsheet QFT of the string in $AdS_5 \times S^5$
- The wrapping graphs contributing at 4-loops correspond to a *single* 'virtual' graph:




- This corresponds to a simple expression

$$\Delta_w^{(4-loop)} = - \sum_{Q=1}^{\infty} \int_{-\infty}^{\infty} \frac{dq}{2\pi} Y_Q(q)$$

where $Y_Q(q)$ is a relatively simple rational function of q and Q

- The Konishi operator corresponds to a two particle state in the 2D worldsheet QFT of the string in $AdS_5 \times S^5$
- The wrapping graphs contributing at 4-loops correspond to a *single* 'virtual' graph:



- This corresponds to a simple expression

$$\Delta_w^{(4-loop)} = - \sum_{Q=1}^{\infty} \int_{-\infty}^{\infty} \frac{dq}{2\pi} Y_Q(q)$$

where $Y_Q(q)$ is a relatively simple rational function of q and Q

- The integral can be easily done by residues giving a remaining sum

$$\sum_{Q=1}^{\infty} \left\{ -\frac{\text{num}(Q)}{(9Q^4 - 3Q^2 + 1)^4 (27Q^6 - 27Q^4 + 36Q^2 + 16)} + \frac{864}{Q^3} - \frac{1440}{Q^5} \right\}$$

where

$$\begin{aligned} \text{num}(Q) = & 7776Q(19683Q^{18} - 78732Q^{16} + 150903Q^{14} - 134865Q^{12} + \\ & + 1458Q^{10} + 48357Q^8 - 13311Q^6 - 1053Q^4 + 369Q^2 - 10) \end{aligned}$$

- This can be summed up giving

$$\Delta_{\text{wrapping}} E = (324 + 864\zeta(3) - 1440\zeta(5))g^8$$

which **exactly agrees** with the direct 4-loop perturbative computation of [F.Fiamberti, A.Santambrogio, C.Sieg and D.Zanon]

- The integral can be easily done by residues giving a remaining sum

$$\sum_{Q=1}^{\infty} \left\{ -\frac{\text{num}(Q)}{(9Q^4 - 3Q^2 + 1)^4 (27Q^6 - 27Q^4 + 36Q^2 + 16)} + \frac{864}{Q^3} - \frac{1440}{Q^5} \right\}$$

where

$$\begin{aligned} \text{num}(Q) = & 7776Q(19683Q^{18} - 78732Q^{16} + 150903Q^{14} - 134865Q^{12} + \\ & + 1458Q^{10} + 48357Q^8 - 13311Q^6 - 1053Q^4 + 369Q^2 - 10) \end{aligned}$$

- This can be summed up giving

$$\Delta_{\text{wrapping}} E = (324 + 864\zeta(3) - 1440\zeta(5))g^8$$

which **exactly agrees** with the direct 4-loop perturbative computation of [F.Fiamberti, A.Santambrogio, C.Sieg and D.Zanon]

- The integral can be easily done by residues giving a remaining sum

$$\sum_{Q=1}^{\infty} \left\{ -\frac{\text{num}(Q)}{(9Q^4 - 3Q^2 + 1)^4 (27Q^6 - 27Q^4 + 36Q^2 + 16)} + \frac{864}{Q^3} - \frac{1440}{Q^5} \right\}$$

where

$$\begin{aligned} \text{num}(Q) = & 7776Q(19683Q^{18} - 78732Q^{16} + 150903Q^{14} - 134865Q^{12} + \\ & + 1458Q^{10} + 48357Q^8 - 13311Q^6 - 1053Q^4 + 369Q^2 - 10) \end{aligned}$$

- This can be summed up giving

$$\Delta_{\text{wrapping}} E = (324 + 864\zeta(3) - 1440\zeta(5))g^8$$

which **exactly agrees** with the direct 4-loop perturbative computation of [F.Fiamberti, A.Santambrogio, C.Sieg and D.Zanon]

- The integral can be easily done by residues giving a remaining sum

$$\sum_{Q=1}^{\infty} \left\{ -\frac{\text{num}(Q)}{(9Q^4 - 3Q^2 + 1)^4 (27Q^6 - 27Q^4 + 36Q^2 + 16)} + \frac{864}{Q^3} - \frac{1440}{Q^5} \right\}$$

where

$$\begin{aligned} \text{num}(Q) = & 7776Q(19683Q^{18} - 78732Q^{16} + 150903Q^{14} - 134865Q^{12} + \\ & + 1458Q^{10} + 48357Q^8 - 13311Q^6 - 1053Q^4 + 369Q^2 - 10) \end{aligned}$$

- This can be summed up giving

$$\Delta_{\text{wrapping}} E = (324 + 864\zeta(3) - 1440\zeta(5))g^8$$

which **exactly agrees** with the direct 4-loop perturbative computation of [F.Fiamberti, A.Santambrogio, C.Sieg and D.Zanon]

- We extended the string computation to give the result up to 5 loops (no gauge theoretical computation so far [Bajnok,Hegedus,RJ,Łukowski])

$$\Delta = 4 + 12 g^2 - 48 g^4 + 336 g^6 + 96(-26 + 6 \zeta(3) - 15 \zeta(5)) g^8 - 96(-158 - 72 \zeta(3) + 54 \zeta(3)^2 + 90 \zeta(5) - 315 \zeta(7)) g^{10}$$

- This could be extended to twist two operators at 4 loops [Bajnok,RJ,Łukowski]
- Nontrivial relations with BFKL and NLO BFKL equations...

- We extended the string computation to give the result up to 5 loops (no gauge theoretical computation so far) [Bajnok,Hegedus,RJ,Łukowski]

$$\Delta = 4 + 12 g^2 - 48 g^4 + 336 g^6 + 96(-26 + 6 \zeta(3) - 15 \zeta(5)) g^8 - 96(-158 - 72 \zeta(3) + 54 \zeta(3)^2 + 90 \zeta(5) - 315 \zeta(7)) g^{10}$$

- This could be extended to twist two operators at 4 loops [Bajnok,RJ,Łukowski]
- Nontrivial relations with BFKL and NLO BFKL equations...

- We extended the string computation to give the result up to 5 loops (no gauge theoretical computation so far [\[Bajnok,Hegedus,RJ,Łukowski\]](#))

$$\Delta = 4 + 12 g^2 - 48 g^4 + 336 g^6 + 96(-26 + 6 \zeta(3) - 15 \zeta(5)) g^8 - 96(-158 - 72 \zeta(3) + 54 \zeta(3)^2 + 90 \zeta(5) - 315 \zeta(7)) g^{10}$$

- This could be extended to twist two operators at 4 loops [\[Bajnok,RJ,Łukowski\]](#)
- Nontrivial relations with BFKL and NLO BFKL equations...

- We extended the string computation to give the result up to 5 loops (no gauge theoretical computation so far [\[Bajnok,Hegedus,RJ,Łukowski\]](#))

$$\Delta = 4 + 12 g^2 - 48 g^4 + 336 g^6 + 96(-26 + 6 \zeta(3) - 15 \zeta(5)) g^8 - 96(-158 - 72 \zeta(3) + 54 \zeta(3)^2 + 90 \zeta(5) - 315 \zeta(7)) g^{10}$$

- This could be extended to twist two operators at 4 loops [\[Bajnok,RJ,Łukowski\]](#)
- Nontrivial relations with BFKL and NLO BFKL equations...


- We extended the string computation to give the result up to 5 loops (no gauge theoretical computation so far [\[Bajnok,Hegedus,RJ,Łukowski\]](#))

$$\Delta = 4 + 12 g^2 - 48 g^4 + 336 g^6 + 96(-26 + 6 \zeta(3) - 15 \zeta(5)) g^8 - 96(-158 - 72 \zeta(3) + 54 \zeta(3)^2 + 90 \zeta(5) - 315 \zeta(7)) g^{10}$$

- This could be extended to twist two operators at 4 loops [\[Bajnok,RJ,Łukowski\]](#)
- Nontrivial relations with BFKL and NLO BFKL equations...

The Konishi operator from string theory


- Recently several groups proposed (infinte) sets of coupled nonlinear integral equations which should work for *any coupling*
[Arutyunov,Frolov],[Bombardelli,Fioravanti,Tateo],[Gromov,Kazakov,Vieira]
- [Gromov,Kazakov,Vieira] proposed concrete equations for the dimension of the Konishi operator for any coupling



- Still some issues to understand - source terms; possibility of reducing the number of equations to a finite number
- Goal:** (*within reach*) – calculate anomalous dimensions for any coupling!!

The Konishi operator from string theory


- Recently several groups proposed (infinte) sets of coupled nonlinear integral equations which should work for *any coupling*
[Arutyunov,Frolov],[Bombardelli,Fioravanti,Tateo],[Gromov,Kazakov,Vieira]
- [Gromov,Kazakov,Vieira] proposed concrete equations for the dimension of the Konishi operator for any coupling



- Still some issues to understand - source terms; possibility of reducing the number of equations to a finite number
- Goal:** (*within reach*) – calculate anomalous dimensions for any coupling!!

The Konishi operator from string theory

- Recently several groups proposed (infinte) sets of coupled nonlinear integral equations which should work for *any coupling*
[Arutyunov,Frolov],[Bombardelli,Fioravanti,Tateo],[Gromov,Kazakov,Vieira]
- [Gromov,Kazakov,Vieira] proposed concrete equations for the dimension of the Konishi operator for any coupling



- Still some issues to understand - source terms; possibility of reducing the number of equations to a finite number
- Goal:** (*within reach*) – calculate anomalous dimensions for any coupling!!

- The agreement of the Konishi computation with the 4-loop weak coupling perturbative gauge theory result is an extremely nontrivial test of AdS/CFT!
- The result came from a single diagram – in contrast to direct perturbative computations in gauge theory which are much more complex
- This suggests that one can use string theory methods of AdS/CFT as an efficient calculational tool also at *weak coupling*
- The AdS/CFT correspondence allows to use methods of exactly solvable integrable **two-dimensional** QFT's to study the **four-dimensional** supersymmetric $\mathcal{N} = 4$ gauge theory
- We may be very close to the complete solution for the spectrum of the theory (\equiv anomalous dimensions at any coupling)

- The agreement of the Konishi computation with the 4-loop weak coupling perturbative gauge theory result is an extremely nontrivial test of AdS/CFT!
- The result came from a single diagram – in contrast to direct perturbative computations in gauge theory which are much more complex
- This suggests that one can use string theory methods of AdS/CFT as an efficient calculational tool also at *weak coupling*
- The AdS/CFT correspondence allows to use methods of exactly solvable integrable **two-dimensional** QFT's to study the **four-dimensional** supersymmetric $\mathcal{N} = 4$ gauge theory
- We may be very close to the complete solution for the spectrum of the theory (\equiv anomalous dimensions at any coupling)

- The agreement of the Konishi computation with the 4-loop weak coupling perturbative gauge theory result is an extremely nontrivial test of AdS/CFT!
- The result came from a single diagram – in contrast to direct perturbative computations in gauge theory which are much more complex
- This suggests that one can use string theory methods of AdS/CFT as an efficient calculational tool also at *weak coupling*
- The AdS/CFT correspondence allows to use methods of exactly solvable integrable **two-dimensional** QFT's to study the **four-dimensional** supersymmetric $\mathcal{N} = 4$ gauge theory
- We may be very close to the complete solution for the spectrum of the theory (\equiv anomalous dimensions at any coupling)

- The agreement of the Konishi computation with the 4-loop weak coupling perturbative gauge theory result is an extremely nontrivial test of AdS/CFT!
- The result came from a single diagram – in contrast to direct perturbative computations in gauge theory which are much more complex
- This suggests that one can use string theory methods of AdS/CFT as an efficient calculational tool also at *weak coupling*
- The AdS/CFT correspondence allows to use methods of exactly solvable integrable **two-dimensional** QFT's to study the **four-dimensional** supersymmetric $\mathcal{N} = 4$ gauge theory
- We may be very close to the complete solution for the spectrum of the theory (\equiv anomalous dimensions at any coupling)

- The agreement of the Konishi computation with the 4-loop weak coupling perturbative gauge theory result is an extremely nontrivial test of AdS/CFT!
- The result came from a single diagram – in contrast to direct perturbative computations in gauge theory which are much more complex
- This suggests that one can use string theory methods of AdS/CFT as an efficient calculational tool also at *weak coupling*
- The AdS/CFT correspondence allows to use methods of exactly solvable integrable **two-dimensional** QFT's to study the **four-dimensional** supersymmetric $\mathcal{N} = 4$ gauge theory
- We may be very close to the complete solution for the spectrum of the theory (\equiv anomalous dimensions at any coupling)

- The agreement of the Konishi computation with the 4-loop weak coupling perturbative gauge theory result is an extremely nontrivial test of AdS/CFT!
- The result came from a single diagram – in contrast to direct perturbative computations in gauge theory which are much more complex
- This suggests that one can use string theory methods of AdS/CFT as an efficient calculational tool also at *weak coupling*
- The AdS/CFT correspondence allows to use methods of exactly solvable integrable **two-dimensional** QFT's to study the **four-dimensional** supersymmetric $\mathcal{N} = 4$ gauge theory
- We may be very close to the complete solution for the spectrum of the theory (\equiv anomalous dimensions at any coupling)