

# "QCD tests at NA48/2"

<u>EPS\_09</u> <u>Krakow 16.7.2009</u>

<u>Gianluca Lamanna</u> Scuola Normale Superiore and INFN Pisa

On behalf of the <u>NA48/2 collaboration</u>: Cambridge, CERN, Chicago, Dubna, Edimburgh, Ferrara, Florence, Mainz, Northwestern, Perugia, Pisa, Saclay, Siegen, Turin, Vienna

# Outline

- NA48/2 experimental setup
- QCD tests
  - ππ scattering lengths (Ke4 and Cusp) (preliminary!)
  - $K^{\pm} \rightarrow \pi^{\pm} \gamma \gamma$  : BR (preliminary!)
  - K<sup>±</sup>→π<sup>±</sup>γe<sup>+</sup>e<sup>-</sup> : BR and Shape (final!)
- Conclusions







- fixed target experiment at CERN-SPS
- Simultaneous, unseparated, focused beams
- 60±3 GeV/c kaon momentum (~7x10<sup>11</sup> ppp)
- 6.3 x 10<sup>7</sup> particles per pulse in decay region
- Similar acceptance for K+ and K- decays
- K+/K- ~ 1.8

#### Detector



- ~100 m long decay region in vacuum
- Triggers based on LKr peaks, CHOD hits and DCH multiplicity
- Similar acceptance between K+ and Kbeams checked reversing magnetics fields
- Pion decay products, from the hadronic beam, remain into the beam pipe

#### Spectrometer:

 $\sigma_p/p = 1.0\% + 0.044\% p$  [p in GeV/c]

#### LKR calorimeter:

 $\sigma_{\rm E}/{\rm E}$  = 3.2%/ $\sqrt{\rm E}$  + 9%/E + 0.42% [E in GeV]

CHOD, HAC, MUV, vetos

#### Kabes

#### **Beam Monitor**



# **Data Taking**



Detector and trigger optimized to collect 3 pions events

Before NA48/2 NA48/2

$$K^{\pm} \rightarrow \pi^{\pm} \pi^{0} \pi^{0}$$
: ~1.10<sup>8</sup>

- Unprecedented statistics in many channels
- Two years of data taking (2003+2004)

• Main purpose was to measure direct CP violation in charged Kaon decays, through asymmetry in Dalitz plot distribution.

New limits on CP violation in charged kaon decays

 $A_g = (-1.5 + 1.5_{stat} + 0.9_{trig} + 1.1_{syst}) \cdot 10^{-4}$ 

 $A_{g}^{0} = (1.8 + 1.7_{stat} + 0.5_{syst}) \cdot 10^{-4}$ 

Phys.Lett.B 634:474 482,2006 Phys.Lett.B 638:22-29,2006 Eur.Phys.J C52:875-891,2007



## **Ke4: Formalism**

• The Ke4 (K<sup>±</sup> $\rightarrow \pi^{+}\pi^{-} e^{\pm}v$ ) dynamics is fully described by 5 (*Cabibbo-Maksymovicz*) variables: M<sub> $\pi\pi^{2}$ </sub>, M<sub>ev</sub><sup>2</sup>, cos $\theta_{\pi}$ , cos $\theta_{e}$  and  $\phi$ 

In the partial waves expansion the amplitude can be written using 2 axial and 1 vector form factors (the axial form factor R is suppressed in  $K_{e4}$  but accessible in  $K_{\mu4}$ ):

$$F = F_{s}e^{i\delta s} + F_{p}e^{i\delta p}\cos\theta_{\pi}$$
$$G = G_{p}e^{i\delta p}$$
$$H = H_{e}e^{i\delta p}$$

F (F<sub>p</sub>,F<sub>s</sub>), G, H and  $\delta = \delta_p - \delta_s$  will be used as fit parameters



• The  $q^2 = (M_{\pi\pi}^2/4m_{\pi}^2)$ -1 dependence can be studied expanding the fitted form factors assuming isospin symmetry:

$$F_{s} = f_{s} + f_{s}'q^{2} + f_{s}''q^{4} + f_{e}'(M_{ev}^{2}/4m_{\pi}^{2}) + \dots$$

$$F_{p} = f_{p} + f_{p}'q^{2} + \dots$$

$$G_{p} = g_{p} + g_{p}'q^{2} + \dots$$

$$H_{p} = h_{p} + h_{p}'q^{2} + \dots$$

## **Ke4: Selection**

#### Signal selection:

- 3 tracks on spectrometer, total charge ±1
- $\bullet$  Missing energy and missing Pt, compatible with  $\nu$
- LKr/DCH used for electron PID
- 2 opposite sign pions
- ~1150000 decays

Main background sources:

 $\pi\pi\pi + \pi \rightarrow ev$ 

 $\pi\pi\pi$  with  $\pi$  misidentified

 $\pi\pi^0\pi^0$  or  $\pi^0 + \pi^0$ (Dalitz) +e misidentified and  $\gamma$ s outside the LKr



The background is studied using the electron *"wrong"* sign (WS) events (we assume  $\Delta Q = \Delta S$  and total charge ±1) and cross check with MC. The total bkg is at level of 0.5%.

7

# Ke4: Fitting procedure



10(Mππ)x5(Mev)x5(cosθe)x5(cosθπ)x12( $\phi$ )=1500 iso-populated boxes

• In each  $M\pi\pi$  bin the F,G,H and  $\delta$  form factors are extracted minimizing a log-likelihood estimator over the 1500 boxes

• All the form factors are measured relatively to fs (no overall BR measured)

• K+ and K- fitted separately, and then the fitted parameters are combined in each  $M\pi\pi$  bin

• The form factors structure is studied in 10 bins of  $q^2$  (or  $M^2_{\phantom{2}\pi\pi})$ 

Agreement with published results (based on 2003 data) [Batley et al. EPJC 54-3 (2008) 411]

(2003+2004)		Data	MC
K+	evts	726400	17.7x10^6
	evts/bin	48	1160
K-	evts	404400	9.8x10^6
	evts/bin	27	650

 $f'_{s}/f_{s} = 0.152\pm0.007\pm0.005$   $f''_{s}/f_{s} = -0.073\pm0.007\pm0.006$   $f'_{e}/f_{s} = 0.068\pm0.006\pm0.007$   $f_{p}/f_{s} = -0.048\pm0.003\pm0.004$   $g_{p}/f_{s} = 0.868\pm0.010\pm0.010$   $g'p/f_{s} = 0.089\pm0.017\pm0.013$   $h_{p}/f_{s} = -0.398\pm0.015\pm0.008$ (stat+syst error guoted)

#### Ke4: Phase shift and $\pi\pi$ scattering length



• The extraction of pion scattering lengths from the fitted  $\delta = \delta_s - \delta_p$  phase shift needs external theoretical and experimental inputs:

The *Roy equations* provide the relation between δ and a0 and a2 near threshold (1) (2) (3)

• Extrapolating the data from the  $M_{\pi\pi}$ >0.8 GeV it's possible to fit the result in the threshold region (the uncertainty from the experimental data defines the *Universal Band*) • Each point in the plot has been corrected for isospin effect: 10-15 mrad

• The total error in each point is different for each experiment:

#### Geneva-Saclay: 40-50 mrad

E865: 15-20 mrad

NA48/2: 7-8 mrad

- Coulomb correction (Gamow factor) and real photons are included in simulation
- Isospin correction prescription given by Gasser (4) results in 11 to 15 mrad in the fitted  $M\pi\pi$  range

• The isospin correction is slightly greater than the error on each point: can't be ignored!

1)[Ananthanarayan,Colangelo,Gasser,Leutwyler Phys.Rept.353:207-279 (2001)] 2)[Descotes-Genon, Fuchs, Girlanda,Stern Eur.Phys.J.C24:469-483,2002] 3)[Kaminski, Pelaez, Yndurain Phys.Rev.D77 (2008)] 4)[Gasser et al. Eur.Phys.J. C59:777,2009]

#### Ke4: a0 and a2 results



## **Cusp: Experimental evidence**

• In  $K^{\pm} \rightarrow \pi^{\pm} \pi^{0} \pi^{0}$  decay the matrix element is usually given as polynominal expansion as a function of the Dalitz variables U and V

 $|M(u,v)|^2 \sim 1 + gu + hu^2 + kv^2 + ...$  $u = (s_3 - s_0)/m_{\pi}^2 \quad s_i = (P_K - P_{\pi,i})^2$ 

•Thanks to the big statistics collected by NA48/2 and the good energy resolution, for the first time a structure has been observed at the  $\pi\pi$  threshold value [Batley et al.,Phys.Lett. B633:173-283,2006]

• This structure has been interpreted by Cabibbo [Cabibbo Phys. Rev. Lett. 93, 121801 (2004)] as due to the strong  $\pi\pi$  rescattering in the K<sup>±</sup> $\rightarrow\pi^{\pm}\pi^{+}\pi^{-}$  final state







## Cusp: $K \rightarrow 3\pi$ selection

Online selection:

• (L1) one charged particle in CHOD and at least three peaks in at least one projection (x or y) of the LKr analog sum peak system.

- (L2) The opposite mass with respect to the charged tracks have to be far from one  $\pi^0$  mass.
- <u>Offline selection</u>: among all the possible  $\gamma$  pairings, the couple for which  $\chi^2$  is smallest is selected
  - the neutral vertex is obtained by imposing the  $\pi^0$  mass. The final vertex is obtained as average between the 2 vertices.





• Very good acceptance and resolution in threshold region: high statistics and good opportunity to resolve tiny structures.

# Cusp: Theoretical approach (Cabibbo-Isidori)

- Fenomenological approach
  The term M0 (no rescattering) is given by the standard PDG expansion
  the first rescattering terms is real below threshold and immaginary above. Interference below threshold.
- $s_{\pi\pi} > (2m_{\pi+})^{2} \qquad M^{2} = (M_{0})^{2} + |M_{1}|^{2}$   $s_{\pi\pi} < (2m\pi_{+})^{2} \qquad M^{2} = (M_{0})^{2} + (M_{1})^{2} + 2M_{0}M_{1}$   $\mathcal{M}_{1} = -2/3(a_{0} a_{2})m_{+}\mathcal{M}_{+} \qquad \sqrt{1 (\frac{M_{00}}{2m_{+}})^{2}}$   $\frac{\pi}{2m_{+}}$

Cabibbo, Isidori JHEP 0503 (2005) 21



- The amplitude below threshold depends on (a0-a2)
- Other 5 terms rise from two loops calculation (proportional to scattering lengths): effects below and above threshold
  Theoretical error evaluated from the next level expansion 13 (work in progress...)

# **Cusp: theoretical approach (Bern-Bonn)**

 Different approach based on effective non-relativistic Lagrangian

- The electromagnetic effects are naturally included in this approach (explicitly omitted in the CI work)
- Different structure of the expansion (different correlation between the terms wrt the Cabibbo-Isidori expansion): kinetic energy and threshold parameter.
- Simultaneus fitting of charged and neutral amplitude to exctract M+ slope parameters (modified with respect to the PDG parametrization)
- Radiative correction, outside the cusp point, included in the BB model



$$\mathcal{M}^{1-\text{loop}} = B_1 J_{+-}(s_3) + B_2 J_{00}(s_3) \\ + \left[ B_3 J_{+0}(s_1) + (s_1 \leftrightarrow s_2) \right]$$

$$\mathcal{M}^{2-\text{loop}} = 2G_0 C_x^2 \underbrace{J_{+-}(s_3) J_{00}(s_3)}_{\text{double loops}} + \dots$$

+ 
$$4H_0C_xC_{+-}$$
  $\underbrace{F_{+}(\ldots;s_3)}_{\text{overlapping loops}}$ 

 $\vdash \ \mathcal{O}(i \, \epsilon^4) \quad [\not\propto \ \text{scatt. lengths}]$ 

[Colangelo, Gasser, Kubis, Rusetsky in Phys.Lett.B638:187-194,2006]

[Bissenger, Fuhrer, Gasser, Kubis, Rusetsky in Phys.Lett.B659:576,2008]

[Bissegger, Fuhrer, Gasser, Kubis, Rusetsky in NPH B806:178, 2009]

# **Cusp: fitting procedure**

 Resolution and detector response matrix obtained using accurate Geant3 based simulation

• Both theories can be fitted with the same procedure (fit parameters: g, h,  $a_0$ - $a_2$ ,  $a_2$ , N)

•The M+ terms appearing in the **CI** theory is fixed by the recent measured  $K^{\pm} \rightarrow \pi^{\pm} \pi^{+} \pi^{-}$  slope parameters [Batley et al. Phys.Lett.B649:349-358,2007]

• In the **BB** the M+ term is obtained simultaneously fitting  $K^{\pm} \rightarrow \pi^{\pm} \pi^{0} \pi^{0}$  and  $K^{\pm} \rightarrow \pi^{\pm} \pi^{+} \pi^{-}$  Dalitz plot

Isospin effects included

• The excess of events in the cusp position, R=(1.8±0.3)·10<sup>-5</sup>, can be explain by:

•pionium 0.8•10<sup>-5</sup> [Silagadze, JETP Lett. 60 (1994) 689]

unbound state with resonant structure
[Gevorkian, Tarasov,
Voskeresenskaya, Phys.Lett.B649:159,2007]



• The numbers of "atoms" is used as free parameter including the 7 bins around the cusp.

#### **Cusp: Results**

2 par. Fit :

 $(a_0-a_2)m_{+}= 0.2571 \pm 0.0048_{\text{stat.}} \pm 0.0025_{\text{syst.}} \pm 0.0014_{\text{ext.}}$  $a_2m_{+}= -0.024 \pm 0.013_{\text{stat.}} \pm 0.009_{\text{syst.}} \pm 0.002_{\text{ext.}}$ 

1 par. Fit (ChPT): (a0-a2)m+=  $0.2633 \pm 0.0024$ stat.  $\pm 0.0014$ syst.  $\pm 0.0019$ ext.



- Results based on BB model (better  $\chi 2$ , most complete theory)
- Very good agreement between the two approches
- Good agreement with ChPT prediction [Colangelo, Gasser, Leutwyler PRL 86 (2001) 5008]:

#### (a0-a2)m<sub>+</sub>=0.265±0.004

- Similar theoretical and experimental precision
- Systematics error dominated by LKr non linearity and trigger efficiency

## **Cusp and Ke4: comparisons**



Very good agreement with  $(a0-a2)m_{+}=0.265 \pm 0.004$ 

ChPT prediction:

 $(a0 - a2)m_{+} = 0.2639 \pm 0.0020$  $\pm 0.0004 \pm 0.0021$ 

Using ChPT constraints:  $(a0 - a2)m_{+} = 0.2640 \pm 0.0020$ ±0.0017 ±0.0035

# K→πγγ: Theory

• At leading order, in the framework of ChPT, the  $K(p) \rightarrow \pi(p1)\gamma(q2)\gamma(q3)$  differential rate is given by:

ifferential rate is given by:  

$$\frac{\partial^2 \Gamma}{\partial y \partial z} = \frac{m_K}{(8\pi)^3} \left[ z^2 (|A+B|^2 + |C|^2) + (y^2 - 1/4\lambda(1,z,r_\pi^2))^2 (|B|^2 + |D|^2) \right]$$

 $y=p(q_2-q_3)/m_{K}^2$  $z=(q_2+q_3)/m_{K}^2$ 

• At O(p4) the relevant contribution is given by  $A(z, \hat{c})$  (loops) and C(z) (poles); B=D=0 at O(p4) and relevant at leading order only at low myy mass (1).

**K(**p)

- ĉ is O(1) and have to be extracted from data
- Order O(p6) can be relevant (30%-40% enhanced in BR) (2).

(1) [Ecker, Pich, De Rafael, in Nucl.Phys.B303:665,1988] (2) [D'Ambrosio, Portoles, in Nucl.Phys.B386:403,1996]

rlp1)

<u>γ(q2)</u>



16.7.2009

## K→πγγ: Results



- Analysis based on 40% of whole data set
- 1164 events found
- main background:  $K^+ \rightarrow \pi^+ \pi^0 \gamma$  ( 3.3%)
- Main systematics from trigger efficency determination
- Data shape (mγγ) follows ChPT description (ĉ=2 shown for qualitative comparison)

 Assuming O(p6) and c=2 the preliminary result on model dependent
 BR determination is

Br(K<sup>+</sup> $\rightarrow \pi^+ \gamma \gamma$ ) = (1.07±0.04<sub>stat</sub> ±0.08<sub>syst</sub>)·10<sup>-6</sup>

(model independent analysis on going)

 Overtaken the previous result from E787 with 31 events candidates:

Br(K<sup>+</sup> $\rightarrow \pi^+ \gamma \gamma$ ) = (1.10±0.32)·10<sup>-6</sup>

## K→πγee: Theory

- Similar with respect to  $K^+ \rightarrow \pi^+ \gamma \gamma$
- Br and m<sub>eeγ</sub> at order O(p4) determined by <sup>C</sup>
- O(p6) could enhance the BR by 30%-40%
- Theoretical prediction at O(p6):

Br(K<sup>+</sup> $\rightarrow \pi^+ \gamma$  e+e-) = (0.9÷1.6)·10<sup>-8</sup>

[Gabbiani, in Phys.Rev.Lett.D59,1999]



• Never observed before!



#### K→πγee : Results



## Conclusions

• The kaon decays give the possibility to study the low energy hadronic interaction with good precision

- Thanks to high statistics and high data quality, NA48/2 can check several ChPT predictions with very high accuracy
- Three significative *"examples"* have been presented in this talk:
  - ππ scattering lengths
  - $K^{\pm} \rightarrow \pi^{\pm} \gamma \gamma$
  - K<sup>±</sup>→π<sup>±</sup>γe<sup>+</sup>e<sup>-</sup>
- The results obtained are compatible with the predictions

• In particular the measurement of the  $\pi\pi$  scattering lengths obtained in two different ways, with an experimental error comparable with the theoretical uncertainty, is a very strong test of the theory.