


Searches for Excited Fermions at HERA

-- Final Results from H1

Z. Zhang (LAL, Orsay) on behalf of the H1 Collaboration



Introduction

➤ HERA:

- Unique $e^{\pm}p$ collider of the world (1992-2007)
- High c.m. energy up to 320GeV for direct searches
- Integrated luminosity/experiment: $\sim 0.5 \text{ fb}^{-1}$

➤ Excited fermions (ν^* , e^* and q^*)

Very actively searched for at HERA (& elsewhere):

8 (H1) + 4 (ZEUS) publications

of which 3 from H1 based on full HERA data

➔ subject of this talk

Motivation & the Model

- 3 family & mass hierarchy \leftrightarrow Composite Model of fermions

↙
Excited fermions

- The effective Lagrangian:

$$\mathcal{L}_{GM} = \frac{1}{2\Lambda} \bar{F}_R^* \sigma^{\mu\nu} \left[\underset{\text{SU}(2)}{g f \frac{\tau^a}{2} W_{\mu\nu}^a} + \underset{\text{U}(1)}{g' f' \frac{Y}{2} B_{\mu\nu}} + \underset{\text{SU}(3)}{g_s f_s \frac{\lambda^a}{2} G_{\mu\nu}^a} \right] F_L$$




Λ : Compositeness scale

f, f', f_s : relative strength to $Z/W, \gamma, g$

Hagiwara et al., Z. Phys. C 29 (1985) 115; Boudjema et al., Z. Phys. C 57 (1993) 425

Productions and Decays

Excited fermions of 1st generation: singly produced at HERA



Decays to standard fermions and gauge bosons
with $Z \rightarrow ee, \mu\mu, qq$, and $W \rightarrow e\nu_e, \mu\nu_\mu, qq'$

→ Experimental signature:
isolated lepton(s), γ , missing energy + jets

Excited Neutrinos

H1 Collab., Phys. Lett. B 663 (2008) 382


□ Use all HERA-2 e-p data (184 pb⁻¹)

(cross section much larger than in e⁺p data)

□ Model Assumption:

$f = -f'$ or $f = +f'$ (insensitive to f_s)

~90%, 85% decay modes investigated





Channel	Data	SM	Signal efficiency [%]
$\nu^* \rightarrow \nu \gamma$	7	12.3 ± 3.0	50–55
$\nu^* \rightarrow eW \rightarrow eq\bar{q}$	220	223 ± 47	40–65
$\nu^* \rightarrow eW \rightarrow e\nu\mu$	0	0.40 ± 0.05	35
$\nu^* \rightarrow eW \rightarrow eve$	0	0.7 ± 0.1	45
$\nu^* \rightarrow \nu Z \rightarrow \nu q\bar{q}$	89	95 ± 21	25–55
$\nu^* \rightarrow \nu Z \rightarrow \nu ee$	0	0.19 ± 0.05	45

→ No evidence for ν^* found, mass dependent exclusion limits derived

Excited Neutrinos (continued)

H1 Collab., Phys. Lett. B 663 (2008) 382

Limits at 95% CL and all channels combined



- If $f/\Lambda=1/M_{\nu^*}$, $M_{\nu^*}<213$ excluded
- Best sensitivity achieved for mass beyond the LEP reach

- HERA1 limits improved by a factor of 3-4
(scan in f'/f)

Excited Electrons

H1 Collab., Phys. Lett. B 666 (2008) 131


□ Use full HERA data (475 pb⁻¹)

(Four-fold increase in stat vs. HERA-1)

□ Model Assumption:

$f=+f'$ (cross section small for $f=-f'$)

80-90% decay modes investigated





Channel	Data	SM	Signal efficiency [%]
$e^* \rightarrow e\gamma$ (ela.)	42	48 ± 4	60–70
$e^* \rightarrow e\gamma$ (inel.)	65	65 ± 8	60–70
$e^* \rightarrow \nu W \rightarrow \nu q\bar{q}$	129	133 ± 32	20–55
$e^* \rightarrow \nu W \rightarrow \nu e\nu$	4	4.5 ± 0.7	60
$e^* \rightarrow eZ \rightarrow e\nu\nu$	4	4.5 ± 0.7	35
$e^* \rightarrow eZ \rightarrow eq\bar{q}$	286	277 ± 62	20–55
$e^* \rightarrow eZ \rightarrow eee$	0	0.72 ± 0.06	60
$e^* \rightarrow eZ \rightarrow e\mu\mu$	0	0.52 ± 0.05	40–15

→ No evidence for e^* found, mass dependent exclusion limits derived

Excited Electrons (continued)

H1 Collab., Phys. Lett. B 666 (2008) 131

Limits at 95% CL and all channels combined



- If $f / \Lambda = 1 / M_{e^*}$, $M_{e^*} < 272$ excluded
- The results extend previously excluded domain at LEP, Tevatron

- GM: Gauge Mediated interactions
- CI: Contact Interactions
- GM interactions dominate

Excited Quarks

H1 Collab., to appear in Phys. Lett. B (June 2009)


- Use full HERA data (475 pb⁻¹)
(12-fold increase in stat vs. prev. results)

- Model Assumption:

$$f = +f' \quad (f_s = 0 \text{ or } f_s \neq 0)$$

s, *u* channels + interference

(previously only *s*-channel with NWA considered)





Channel	Data	SM	Signal Efficiency [%]
$q^* \rightarrow q\gamma$	44	$46 \pm 4 \pm 7$	35 – 45
$q^* \rightarrow qW/Z \rightarrow qq\bar{q}$	341	$326 \pm 48 \pm 62$	5 – 55
$q^* \rightarrow qW \rightarrow qe\nu$	6	$6.0 \pm 0.2 \pm 0.8$	20 – 30
$q^* \rightarrow qW \rightarrow q\mu\nu$	5	$4.4 \pm 0.2 \pm 0.7$	20 – 40
$q^* \rightarrow qZ \rightarrow qee$	0	$0.44 \pm 0.06 \pm 0.04$	15 – 30
$q^* \rightarrow qZ \rightarrow q\mu\mu$	0	$0.87 \pm 0.10 \pm 0.04$	15 – 30

→ No evidence for q^* found, mass dependent exclusion limits derived

Excited Quarks (continued)

H1 Collab., to appear in Phys. Lett. B (June 2009)

Limits at 95% CL and all channels combined



→ If $f/\Lambda = 1/M_{q^*}$, $M_{q^*} < 252$ excluded

→ The results extend

- beyond the kinematical limits
- previously excluded domain at LEP

→ HERA limits complementary to that of Tevatron for $f_s < f, f'$

Summary

- H1 has given the final word on the searches for f^* :
If $f/\Lambda=1/M_{f^*}$, $M_{f^*}<213, 272, 252\text{GeV}$ excluded at 95%CL
for ν^* , e^* , and q^* , respectively
- The HERA limits with the full data sample
 - improve significantly previous HERA results
 - extend/compete/complement those at LEP/Tevatron