

New physics sensitivity of the rare decay mode

$$B \rightarrow K^0 \ell^- e \ell^+$$

Friday, July 17, 2009 9:30 AM (15 minutes)

We design new observables available from the angular distributions of the decay

$B \rightarrow K^0 e \ell^- e \ell^+$ with high sensitivity to specific new physics operators.

We present a NLO analysis of all observables based on the QCD factorization approach in the low-dilepton mass region and make the uncertainties due to the Λ/m_b corrections manifest and finally analyse the sensitivity of the new observables to new physics.

We explore the experimental sensitivities at LHCb and SuperLHCb based on a full-angular fit method. We also show that the previously discussed transversity amplitude AT_1 cannot be measured at the LHCb experiment or at future B factory experiments as it requires a measurement of the spin of the final state particles.

We also analyse CP violating observables in this decay mode and critically discuss their sensitivity to new physics.

Primary author: Dr HURTH, Tobias (CERN)

Co-authors: Dr MATIAS, Joaquim (Universitat Autònoma de Barcelona); Mr RAMON, Marc (Universitat Autònoma de Barcelona); Dr EGEDE, Ulrik (Imperial College, LHCb); Mr REECE, Will (Imperial College, LHCb)

Presenter: Dr HURTH, Tobias (CERN)

Session Classification: II. Flavour Physics

Track Classification: Flavour Physics