Multiscale phenomena in molecular matter



Contribution ID: 80

Type: poster presentation

## How does the molecular structure of the bent-core molecules affects the occurrence of twist-bend nematic phase?: a mean-field study

Tuesday, 4 July 2017 19:10 (2 hours)

A newly observed –heliconical (twist–bend) nematic phase (NTB) of nanoscale pitch [1] has recently caught much of attention. Initially, the theoretical concept of this phase has been presented by R. B. Meyer [2] in 1973, where he assumed that the director of the molecules precesses on a cone forming an oblique helicoidal structure. Subsequently in 2001 Dozov [3], in 2002 Memmer [4] and in 2004 Lorman et al. [5], suggested that the formation of the N<sub>TB</sub> can be facilitated by the shape of bent–core molecules.

Here we present theoretical investigations on relation between biaxiality and bend angle on stability of nematic twist-bend phase. We use a generalized Maier-Saupe type of model for NTB [6], with intrinsic molecular biaxiality on both of the arms of the molecule, quantified by a  $3 \times 3$  second-rank traceless symmetric tensor **Q**. Values of bend angle ranging from 130 to 170 degree were investigated. The detailed results show the influence of the molecular biaxiality parameter [7] on the relative stability of the NTB, nematic, isotropic phase and the related phase transitions. For the bend angles equal or larger than 140 degree we observe all five possible phases: uniaxial and biaxial twist-bend nematics, together with uniaxial and biaxial nematics and isotropic phase.

Acknowledgments

This work was supported by Grant No. DEC–2013/11/B/ST3/04247 of the National Science Centre in Poland and in part by PL–Grid Infrastructure. W. T. acknowledges support from Marian Smoluchowski Scholarship (KNOW/58/SS/WT/2016) granted by Marian Smoluchowski Cracow Scientific Consortium "Matter–Energy–Future" within the KNOW funding.

## References

[1] V. Borshch, Y-K Kim, J. Xiang, M. Gao, A. Jakli, V. P. Panov, J. K. Vij, C. T. Imrie, M-G. Tamba, G. H. Mehl and O. D. Lavrentovich, *Nature Communications* 4, 2635 (2013).

[2] R. B. Meyer, "Structural Problems in Liquid Crystal Physics", pp. 273-373 in Les Houches Summer School in Theoretical Physics, 1973. Molecular Fluids, R. Balian and G. Weil (eds.) (Gordon and Breach, New York, 1976).

- [3] I. Dozov, EPL (Europhysics Letters) 56, 247 (2001).
- [4] R. Memmer, Liq. Cryst. 29, 483 (2002).
- [5] V. L. Lorman and B. Mettout, *Phys. Rev. E* 69, 061710 (2004).
- [6] W. Tomczyk, G. Pająk, and L. Longa, Soft Matter 12, 7445 (2016).

[7] R. Berardi, L. Muccioli, S. Orlandi, M. Ricci, and C. Zannoni, J. Phys.: Condens. Matter 20, 463101.1 (2008).

**Primary author:** Mr TOMCZYK, Wojciech (Marian Smoluchowski Institute of Physics, Jagiellonian University, prof. S. Łojasiewicza 11 Street, 30-348 Kraków)

**Co-authors:** Dr PAJĄK, Grzegorz (Marian Smoluchowski Institute of Physics, Jagiellonian University, prof. S. Łojasiewicza 11 Street, 30-348 Kraków); Prof. LONGA, Lech (Marian Smoluchowski Institute of Physics, Jagiellonian University, prof. S. Łojasiewicza 11 Street, 30-348 Kraków)

**Presenter:** Mr TOMCZYK, Wojciech (Marian Smoluchowski Institute of Physics, Jagiellonian University, prof. S. Łojasiewicza 11 Street, 30-348 Kraków)

Session Classification: Poster Session

Track Classification: Soft matter and glass formers